



## Bacterial community assemblages in sediments under high anthropogenic pressure at Ichkeul Lake/Bizerte Lagoon hydrological system, Tunisia

Fida Ben Salem, Olfa Ben Saïd, Cristiana Cravo-Laureau, Ezzeddine Mahmoudi, Noëlle Bru, Mathilde Monperrus, Robert Duran

### ► To cite this version:

Fida Ben Salem, Olfa Ben Saïd, Cristiana Cravo-Laureau, Ezzeddine Mahmoudi, Noëlle Bru, et al.. Bacterial community assemblages in sediments under high anthropogenic pressure at Ichkeul Lake/Bizerte Lagoon hydrological system, Tunisia. *Environmental Pollution*, 2019, 252, pp.644-656. 10.1016/j.envpol.2019.05.146 . hal-02170012

**HAL Id: hal-02170012**

**<https://univ-pau.hal.science/hal-02170012>**

Submitted on 25 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

**Bacterial community assemblages in sediments under high anthropogenic pressure at  
Ichkeul Lake / Bizerte Lagoon hydrological system, Tunisia**

**Fida Ben Salem<sup>a,b,d</sup>, Olfa Ben Said<sup>a,b,d</sup>, Cristiana Cravo-Laureau<sup>b,d</sup>, Ezzeddine  
Mahmoudi<sup>a</sup>, Noëlle Bru<sup>c,d</sup>, Mathilde Monperrus<sup>b,d</sup>, Robert Duran<sup>b,d\*</sup>**

<sup>a</sup>Laboratoire de Biosurveillance de l'Environnement, Faculté des Sciences de Bizerte, Tunisie.

<sup>b</sup>MELODY group, Université de Pau et des Pays de l'Adour, E2S-UPPA, IPREM UMR  
CNRS 5254, BP 1155, Pau Cedex, 64013, France

<sup>c</sup>Laboratoire de Mathématiques et de leurs Applications, PAU UMR CNRS 5142 – Université  
de Pau et des Pays de l'Adour, E2S-UPPA, France

<sup>d</sup>Fédération de recherche MIRA, Université de Pau et des Pays de l'Adour, E2S-UPPA,  
France

\*Corresponding author present address:

Pr. Robert Duran

Université de Pau et des Pays de l'Adour

Bâtiment IBEAS

BP1155

64013 Pau Cedex, France

Tel: +33559407568; Fax: +33559407490

Email: Robert.duran@univ-pau.fr

## 26    **Highlights**

- 27        •    Evidence of bacterial continuum in Ichkeul Lake / Bizerte Lagoon
- 28        •    The lake and the lagoon have specific bacterial assemblages
- 29        •    Each type of pollutant is associated with specific bacterial genera
- 30        •    Uncontaminated sediments have also specific bacterial genera
- 31        •    Potential to develop tools for managing bacterial resources

32

## **Abstract**

Bacterial communities inhabiting sediments in coastal areas endure the effect of strong anthropogenic pressure characterized by the presence of multiple contaminants. Understanding the effect of pollutants on the organization of bacterial communities is of paramount importance in order to unravel bacterial assemblages colonizing specific ecological niches. Here, chemical and molecular approaches were combined to investigate the bacterial communities inhabiting the sediments of the Ichkeul Lake / Bizerte Lagoon, a hydrological system under anthropogenic pressure. Although the microbial community of the Ichkeul Lake sediment was different to that of the Bizerte Lagoon, common bacterial genera were identified suggesting a lake-lagoon continuum probably due to the hydrology of the system exchanging waters according to the season. These genera represent bacterial "generalists" maintaining probably general biogeochemical functions. Linear discriminant analysis effect size (LEfSe) showed significant differential abundance distribution of bacterial genera according to the habitat, the pollution type and level. Further, correlation analyses identified specific bacterial genera which abundance was linked with pesticides concentrations in the lake, while in the lagoon the abundance of specific bacterial genera was found linked with the concentrations of PAHs (Polycyclic aromatic hydrocarbons) and organic forms of Sn. As well, bacterial genera which abundance was not correlated with the concentrations of pollutants were identified in both lake and lagoon. These findings represent valuable information, pointing out specific bacterial genera associated with pollutants, which represent assets for developing bacterial tools for the implementation, the management, and monitoring of bioremediation processes to mitigate the effect of pollutants in aquatic ecosystems.

**Keywords:** 16S rRNA sequencing, pollution, pesticide, PAH, network analysis

## 1. Introduction

Lagoons and lakes are highly productive ecosystems, providing important ecological services such as breeding grounds and nurseries for many fish and bird species (Vasconcelos et al., 2011). They are threatened by human activities, and therefore they need careful monitoring because they are very vulnerable (Pérez-Domínguez et al., 2012). Their status may change rapidly in response to natural- and human-driven events (Cravo-Laureau et al., 2017a; Duran et al., 2015c). Sediments are complex habitats densely colonized by diverse microorganisms (Brito et al., 2009; Chronopoulou et al., 2013; Duran et al., 2015a; Fourcans et al., 2008). In sediments, microbial communities are involved in the decomposition of organic matter (Schultz and Urban, 2008). Microorganisms play a key role in carbon, nitrogen and phosphorus cycling (Andrews et al., 2011) as well as in determining the fate of pollutants (Bordenave et al., 2004a; Bordenave et al., 2004b; Guermouche M'rassi et al., 2015; Pringault et al., 2008; Stauffert et al., 2013). Polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs) and organochlorine pesticides (OCPs) are considered as persistent organic pollutants (POPs), which are ubiquitous in aquatic ecosystems (Doong et al., 2008; Kjellerup et al., 2014; Mwanamoki et al., 2014). POPs are mainly derived from industrial and agricultural sources, as well as maritime traffic (Chang and Doong, 2006; Wen et al., 1994). They enter aquatic environments through atmospheric deposition and run-off (Botta et al., 2009; Duran and Cravo-Laureau, 2016). Their fate in the marine environment depends on abiotic and biotic factors (for review, see Duran and Cravo-Laureau, 2016). The rapid development of molecular methods for microbial ecology has provided detailed information on microbial diversity and dynamics in marine sediments (Bordenave et al., 2008; Stauffert et al., 2014; Terrisse et al., 2017; Thiagarajan et al., 2010). The molecular methods, particularly high-throughput sequencing technologies have been proved to be useful tools for bio-monitoring aquatic ecosystem (Cravo-Laureau and Duran, 2014; Cravo-Laureau et al.,

2017b). Their utilization in combination with chemical analyses might shed light on the ecological role and the adaptation of bacteria to environmental factors. Correlation analyses allowed the exploration of the interactions between microorganisms and their habitat (Barberan et al., 2012; Fuhrman and Steele, 2008; Llado et al., 2015; Williams et al., 2014; Yergeau et al., 2012). Particularly, such analyses have identified possible pollutant degraders (Llado et al., 2015; Yergeau et al., 2012), thereby obtaining valuable information for guiding the management of microbial communities to mitigate the effect of pollutants.

The Ichkeul Lake / Bizerte Lagoon hydrological system (Tunisia) is located in an area under high anthropic pressure. The Ichkeul Lake, Ramsar protected area, is important for migrating birds (Ramdani et al., 2001). The contaminants entering into the Ichkeul Lake / Bizerte Lagoon hydrological system (Stevenson and Battarbee, 1991; Trabelsi and Driss, 2005) have been shown to affect aquatic organisms and migratory birds (Ben Said et al., 2009; Casagrande and Boudouresque, 2010; Gharbi-Bouraoui et al., 2008). During the past decades several projects have studied the PAHs in the sediment of Bizerte Lagoon and their impact on the benthic microorganisms and organisms (Ben Said et al., 2009; Ben Said et al., 2008; Ben Said et al., 2015; Louati et al., 2013a; Louati et al., 2013b; Trabelsi and Driss, 2005). Recently, we characterized the contamination of the Ichkeul Lake / Bizerte Lagoon hydrological system. Almost all stations exceeded the threshold effect level (TEL) of the standards published by NOAA (Ben Salem et al., 2017). The Bizerte Lagoon has been shown to contain industrial contaminants including hydrocarbon (PAHs, polycyclic aromatic hydrocarbons) and metallic compounds, while the Ichkeul Lake was found to contain agrochemicals contaminants including organochlorine pesticides such as endrin and dieldrin (Ben Salem et al., 2016a; Ben Salem et al., 2016b; Ben Salem et al., 2017). In a previous study, based on 16S rRNA gene T-RFLP (terminal restriction fragment length polymorphism) analysis investigating the bacterial communities in four contrasted stations of the Ichkeul

Lake / Bizerte Lagoon hydrological system (two located at the Ichkeul Lake, and two at the Bizerte Lagoon), we demonstrated that bacterial community compositions were linked with the type and level of pesticide contamination at the dry season (Ben Salem et al., 2016a). The current study aims to characterize the bacterial communities from the sediments of the Ichkeul Lake / Bizerte Lagoon hydrological system. We further characterize the bacterial communities structure by 16S rRNA genes T-RFLP analysis, expanding the study by investigating 20 stations around the Ichkeul Lake / Bizerte Lagoon hydrological system at two contrasted seasons (April, dry season; September, wet season). Also, we further characterize the bacterial composition of four contrasted stations by sequencing 16S rRNA gene PCR-amplified fragments via 454-pyrosequencing. The effect of organic contaminants such as PAHs, PCBs, OCPs on the bacterial assemblages were determined. Specific bacteria, which abundance was linked with the presence of a pollutant, were identified.

## **2. Materials and Methods**

### **2.1. Site description**

The Ichkeul Lake / Bizerte Lagoon hydrological system, North Tunisia, suffers the effects of human activities because it is located in an area under high anthropic pressure. Bizerte Lagoon has a central role because connected to the Mediterranean Sea and to the Ichkeul Lake via the Tinja River; the hydrological system receives water via different rivers (Oueds) all around (Fig. 1). The lake experiences large fluctuations in salinity each year, from freshwater with average salinity around 4 PSU (Practical salinity unit) in winter (wet season) due to rivers flood, to saline ecosystem with average salinity around 35 PSU in summer (dry season) receiving seawater from the Bizerte Lagoon and because of evaporation. In a previous study,

we demonstrated that pollution levels fluctuated according to seasons (Ben Salem et al., 2016a).

## **2.2. Sampling**

Sediment samples were collected from 20 stations (ST) located all around the Ichkeul Lake / Bizerte Lagoon hydrological system (Fig. 1): ST1 to ST8 in the lake and ST11 to ST20 in the lagoon, ST9 on the channel connecting the lagoon to the sea, and ST10 in the Mediterranean Sea. The lake and lagoon stations were located at the river mouths (13 stations), close to industrial and agricultural areas (4 stations), and at the centers of the lake and the lagoon (2 stations). Two sampling campaigns were performed: September 2011 (wet season) and April 2012 (dry season). Surface sediments (0-10 cm) were collected in triplicates (3 independent samples, i.e. biological replicates) at each station with a Van Veen grab as previously described (Ben Salem et al., 2016a). Collected sediments were dispatched in cryotubes (3 technical replicates were taken), quickly frozen in liquid nitrogen for microbial analyses, and in glass bottles kept at 4°C for chemical analyses. At lab, samples for microbial analyses were stored at -80°C and those for chemical analyses at -20°C until use.

## **2.3. Physical-chemical analyses**

Physical-chemical parameters, including temperature, dissolved oxygen, pH, salinity, phosphate, ammonia, nitrite, and nitrate were determined as previously described (Ben Salem et al., 2016a). Chemical analyses for contaminants followed the QuEChERS procedure described in Ben Salem et al. (2016b). Chemical data and physical-chemical parameters are summarized in supplementary material (Table S1). All chemical data and physical-chemical parameters are presented in details in Ben Salem et al. (2017).

The contamination contents of the sediments were compared to sediment quality guidelines (SQG) in order to determine their potential toxic effect. Sediments were classified according the quality criteria developed by Bakke et al. (2010) for PCBs, OCPs, and PAHs. The toxic



potential of the sediments was evaluated by calculating the mean effect range medium (ERM) quotient (m-ERM-Q) as previously described (Long et al., 1998).

#### **2.4. DNA extraction, PCR and T-RFLP analyses**

DNA extractions were performed using 250 mg of frozen sediment with the UltraClean soil DNA isolation Kit (MoBio Laboratories, CA) as previously described (Precigou et al., 2004). The bacterial universal primers 1387R 5'-GGGCGGWGTGTACAAGGC-3' and 63F 5'-CAGGCCTAACACATGCAAGTC-3' (Marchesi et al., 1998) were used for 16S rRNA gene amplifications. Primers were fluorescently labeled for T-RFLP analysis, which was performed as previously described (Precigou et al., 2001) using 10 ng of extracted DNA. PCR mixture and conditions were performed as previously described (Vercraene-Eairmal et al., 2010). PCR amplicon purification, digestion and electrophoresis separation have been previously described (Vitte et al., 2011). T-RFLP profiles were analyzed as previously described (Cravo-Laureau et al., 2011).

#### **2.5. Sequencing and data analysis**

Among the 20 stations, 4 stations were selected according to the environmental parameters for 454 pyrosequencing analyses. The analysis was performed with the samples collected at the dry season (April 2012) because the highest levels of contaminants (particularly PAHs and pesticides) were recorded (supplementary material Table S1), and also because important contrasted conditions were detected between Ichkeul Lake and Bizerte Lagoon (Table 1, and supplementary material, Table S1). Two stations from the Ichkeul Lake (salinity  $2.6 \pm 0$ , pH  $8.29 \pm 0.02$ ; supplementary material Table S1) were chosen for contrasted contaminations: ST5, showing multiple contaminations by PAHs and pesticides; and ST8, the most contaminated by OCPs (Table 1; supplementary material, Table S1). Similarly, two stations from the Bizerte Lagoon (salinity  $29.5 \pm 0.4$ , pH  $8.37 \pm 0.03$ ) were chosen for contrasted

contaminations: ST12, the most contaminated by PAHs; and ST17, the less contaminated of the entire complex (Table 1; supplementary material, Table S1). Three replicates for each station were sequenced at Molecular Research LP DNA (MR DNA sequencing service, Shallowater, TX, USA) using Roche 454 FLX titanium as previously described (Dowd et al., 2008; Duran et al., 2015a).

QIIME (Quantitative Insights Into Microbial Ecology; Caporaso et al., 2010) was used to analyse the 16S rRNA gene sequence reads as described (Terrisse et al., 2017). Briefly, data cleaning was performed according to Quince et al. (2009) by denoising of the data and by eliminating chimera and sequences with length under 450 bp, with mistakes in primers sequences or homopolymers. OTU picking was performed following the Usearch method (Edgar, 2010) defining reference sequences by applying a 97% similarity threshold. The taxonomic assignment was performed with BLASTn against May 2013 Greengenes database (DeSantis et al., 2006; McDonald et al., 2012) to establish the OTU table. The OTU abundances for each sample were rarefied to the minimum read number observed (12,977 sequences) for a sample using the R package vegan (Oksanen et al., 2013; Version: 2.0–10). The complete dataset was deposited in the NCBI Sequence Read Archive (SRA) database (SUB5043721). It is available under the Bioproject ID PRJNA515186.

## **2.6. Statistical analysis**

Canonical Correspondence Analysis (CCA) correlating bacterial community structure (T-RFLP patterns) with chemical data (PAH and pesticide concentrations) and physical-chemical parameters, was performed with MVSP v3.22 software (Kovach Computing Service, Anglesey Wales). T-RFLP profiles were normalized by calculating relative abundances of each TRF, as previously described (Dominique et al., 2007). Data were square root transformed to get a normal distribution of the data (Dollhopf et al., 2001).

All statistical analyses based on sequencing data were performed on the random subsample of 12,977 bacterial sequences, corresponding to the smaller number of sequences per sample in the dataset. Alpha diversity indices (Shannon-Wiener  $H'$  and richness  $R$ ), based on rarefied OTU table, were calculated using PRIMER v5.0 (Plymouth Routines in Multivariate Ecological Research). The overall differences of these indices between the stations were tested by ANOVA and Tukey HSD multiple comparisons (pairwise comparisons). Significant difference was assumed when  $p < 0.05$ . The contribution of individual OTUs explaining the differences between stations was determined by SIMPER (similarity percentages) based on Bray-Curtis distance measure.

Linear discriminant analysis effect size (LEfSe; Segata et al., 2011) was performed to determine biomarkers of contamination and habitat as previously described (Jeanbille et al., 2016b). Briefly, the non-parametric Kruskal-Wallis sum-rank test ( $\alpha = 0.05$ ) was performed to detect taxa with significant differential abundance. The biological consistency was investigated by performing a pairwise Wilcoxon test ( $\alpha = 0.05$ ). A LDA threshold score of 2.0 and 1000 bootstrap iterations were applied. Heatmap and clustered correlation (Weinstein et al., 1997) were performed with CIMminer (<http://discover.nci.nih.gov/cimminer/>). These analyses were performed at the genus level derived from the OTU abundance table in order to determine the correlations between bacterial genera abundances and pollutant concentrations. Correlation analysis was based on Pearson correlation. The hierarchical clustering of correlations was based on ward method. Strong correlations ( $>|0.5|$ ) were considered. The distribution of OTUs identified in the bacterial patterns of the different stations was visualized with Venn diagrams constructed with Venny using OTU occurrence table as previously described (Duran et al., 2015a). OTUs present at least in two of the triplicates were retained for the analysis.

### 3. Results and discussions

#### 3.1. Contamination level

The chemical analyses of the sediments collected during September 2011 and April 2012 sampling campaigns (supplementary material, Table S1) have been presented in detail in Ben Salem et al. (2017). Table 1 summarizes the chemical data showing the sum of the different types of pollutants obtained in April 2012 for four representative and contrasted stations: 2 from the lake (ST5 and ST8) and 2 from the lagoon (ST12 and ST17). The Bizerte Lagoon was characterized by PAH concentrations ranging from  $137 \pm 26$  to  $17988 \pm 1183 \text{ ng g}^{-1}$ , corresponding to class I (ST12, insignificantly polluted) and class IV (ST17, severely polluted), respectively according to the classification developed by Bakke et al. (2010). Both stations of the Ichkeul Lake (ST5 and ST8) were classified as class II with PAH concentrations of  $1362 \pm 560$  and  $444 \pm 24 \text{ ng g}^{-1}$ , respectively. Considering the contamination by organochlorine pesticides (OCPs), the Ichkeul Lake stations were classified as class IV (severely polluted) while the Bizerte Lagoon stations were classified as class III (markedly polluted) according to the Bakke et al. (2010) classification. The OCPs, including endrin, dieldrin and lindane, and their derivatives, were detected in the Ichkeul Lake at concentrations ranging from  $160 \pm 14$  to  $2021 \pm 85 \text{ ng g}^{-1}$ . Additionally, the chemical analyses revealed low PCB contamination of the sediments (ranging from  $2.48 \pm 0.8$  to  $5.58 \pm 1 \text{ ng g}^{-1}$ ). The sediments were classified as class I (insignificantly polluted) except for the station ST8 that was classified as class II (moderately polluted). In order to take into account the multiple contaminations, the mean effect range medium quotients (m-ERM-Q) were calculated (Table 1; supplementary material, Table S1). The m-ERM-Q showed that the toxic potential of the sediments was higher ( $p$  value  $<0.01$ ) in Ichkeul Lake stations than in Bizerte Lagoon stations at both seasons (supplementary material, Table S1). The station were classified according to increasing m-ERM-Q potential toxicity of sediments as follow ST17

< ST12 < ST5 < ST8 (Table 1). Station ST17 was considered as lowly contaminated, while stations ST5, ST8 and ST12 were considered as highly contaminated.

The chemical data and physical-chemical parameters (supplementary material, Table S1) presented in detail in Ben Salem et al. (2017) were used in the present study to characterize the bacterial community structure according to environmental parameters.

### **3.2. Bacterial community structure**

Correspondence canonical analysis (CCA) using T-RFLP patterns and chemical data separated bacterial communities of the two ecosystems, lake and lagoon irrespective of the season (Fig. 2a). Regarding the environmental parameters, the salinity, the temperature, and the concentrations of nitrite and ammonia were the main factors influencing the bacterial community composition (based on T-RF abundances) of the lagoon, while the bacterial community composition of the lake correlated with dissolved oxygen (DO) and phosphate concentrations. This observation was consistent with previous studies showing that bacterial communities are affected by salinity (Jeanbille et al., 2016a; Jeanbille et al., 2016b; Xi et al., 2014) and nutrients (Bodoque et al., 2017; Liu et al., 2015; Yao et al., 2017). Particularly, nitrite and ammonia concentrations may affect the bacterial populations involved in the nitrogen cycle in lagoon sediments as shown for saline lake sediments (Lipsewiers et al., 2016) and marine contaminated sediments (Mahmoudi et al., 2015). The bacterial populations involved in the nitrogen cycle have been shown to play an important role in hydrocarbon removal in marine sediments (Mahmoudi et al., 2015; Stauffert et al., 2016). The CCA also showed that the pH and the concentrations of nitrate and organo-tin compounds influenced slightly the bacterial community compositions probably because they varied within a small range in the studied sediments.

Regarding the contaminants, the CCA analysis showed that the bacterial community composition was correlated with pesticide concentrations in the lake, while bacterial

community composition was linked with PAHs concentrations in the lagoon (Fig. 2a). These observations confirmed those previously reported (Ben Said et al., 2009) that showed the impact of PAHs on bacteria communities in Bizerte Lagoon sediments. Considering the lake, the bacterial community composition was correlated with PAHs concentrations in April, except for those of station ST8 that was correlated with PCBs and Pesticides concentrations, while bacterial community composition was correlated with the concentration of organo-tin compounds in September (Fig. 2b). This observation suggested seasonal fluctuations in the input of contaminants. It is likely that the lake receives pesticide inputs from crop treatments in April. Additionally, the hydrology of the ecosystem might also play a crucial role: during the dry season (April) lagoon water contaminated by PAHs enters into the lake (Ben Salem et al., 2017). Considering the lagoon, the bacterial community composition was correlated with PAHs and PCBs concentrations in April (Fig. 2c), suggesting that the pollutants were concentrated (concentrating effect) during the dry season. It is likely that the dry season represent a critical season for the Ichkeul Lake / Bizerte Lagoon hydrological system because the environmental conditions favor the concentrating effect of pollutants in both the lake and the lagoon. It is thus of paramount importance to investigate how benthic bacterial communities are affected by such inputs of pollutants. The sediments from the stations ST5 and ST8 (lake) and the stations ST12 and ST17 (lagoon) collected in April were retained for further bacterial characterization because they presented different type of contamination while slightly influenced by environmental parameters (Fig. 2b,c; Table 1).

### **3.3. Bacterial community composition**

In order to further investigate the effect of pollutants on bacterial communities at the dry season (April), the composition of bacterial communities was determined by high throughput pyrosequencing. Four stations showing contrasted chemical characteristics were chosen (Table 1): station ST17 (lagoon), the less contaminated; station ST12 (lagoon), the most

contaminated with PAHs; station ST8 (lake), the most contaminated with OCPs (pesticides); and station ST5 (lake), contaminated with PAHs and OCPs mixture. Three replicates from each station were analysed. After trimming, the 191,278 sequences (400 bp) retained (12,977 sequences per sample after resampling) were dispatched within 858 OTUs, which were grouped into 44 different phyla (Fig. 3). Rarefaction curves (Supplementary data Fig. S1) showed high sequencing coverage in all samples. The coverage (~ 94%) indicated that the sequencing effort was sufficient to assess the bacterial community diversity (Table 2). The sequence affiliation showed that Proteobacteria largely dominated the sediments, with predominance of Deltaproteobacteria (24%), Gammaproteobacteria (23%) and Alphaproteobacteria (9%); Epsilonproteobacteria and Betaproteobacteria accounting for 6% and 5% respectively. The dominance of Proteobacteria, particularly Deltaproteobacteria and Gammaproteobacteria, has been often reported in both lake and lagoon sediments (Borin et al., 2009; Obi et al., 2016; Pavloudi et al., 2016; Quero et al., 2015; Sun et al., 2013) suggesting that they have a pivotal role in biogeochemical cycles in sediment environments. Besides Proteobacteria, Bacteroidetes (9%), Chloroflexi (6%), Planctomycetes (2%) and Nitrospirae (2%) were the major phyla detected in all four stations (Fig. 3), which was consistent with previous observation in sediment environments (Behera et al., 2017; Fan et al., 2016). These phyla have been detected in freshwater and marine sediments (Behera et al., 2017; Fan et al., 2016; Jacquiod et al., 2018b; Paissé et al., 2012; Pavloudi et al., 2016). Bacteroidetes are well known for their capacity to degrade organic matter (Hunter et al., 2006; Zeng et al., 2011). They were in high abundance in contaminated sediments (Paisse et al., 2008; Quero et al., 2015; Sun et al., 2013). Chloroflexi have been found as main players in the transformation of organic compounds including organo-chlorines such as PCBs and pesticides (Fagervold et al., 2007; Hug et al., 2013; Matturro et al., 2016a). Planctomycetes, which have the capability to oxidize ammonia aerobically and anaerobically (Junier et al., 2010), have

been widely detected in contaminated sediments (Dell'Anno et al., 2012; Quero et al., 2015). Similarly, Nitrospirae members playing an important role in the nitrogen cycle (Daims, 2014) have been detected in contaminated sediments by organic compounds including pesticides and PAHs (Ding et al., 2017; Fernanda Sánchez-Soto Jiménez et al., 2018; Stauffert et al., 2013). Alpha diversity indices were not significantly different between stations (Table 2), an observation that was in contradiction with other studies showing that the presence of pollutants either reduce (Haller et al., 2011; Johnston and Roberts, 2009) or increase (Duran et al., 2015a; Jacquiod et al., 2018a) bacterial diversity. It was thus surprising to observe similar bacterial diversity indexes in sediments presenting different pollutant types and levels. Similar observations have been reported by comparing uncontaminated and contaminated sediments (Paisse et al., 2008; Yin et al., 2015) and soils (Berg et al., 2012). The pollution history, particularly the fluctuant and successive pollution loads resulting in legacy effects, may explain the fact that bacterial diversity was maintained (Jurgburg et al., 2017; Nunes et al., 2018). Also, the coalescence of bacterial communities, mixing bacterial communities from different ecosystems (Mansour et al., 2018; Rillig et al., 2015), here lake and lagoon, may contribute to attenuate the diversity differences between the ecosystem, as previously described in river sediments (Yin et al., 2015). The Venn diagram, showing shared OTUs between lake and lagoon stations (Fig. 4a) evidenced the ecological continuum between the lake and the lagoon, which are closely linked via the Tinja River (Fig. 1). Although similar diversity indexes were observed, the composition of the bacterial diversity was different. Indeed, the Venn diagram revealed that each station was characterized by specific OTUs, only 211 OTUs being shared between the stations (Fig. 4a). Such observation was consistent with previous reports on bacterial communities in coastal areas (Duran et al., 2015b; Ikenaga et al., 2010; Misson et al., 2016), indicating that bacterial communities were structured according to the environmental conditions, as revealed by the T-RFLP analyses (Fig. 2). Additionally,



linear discriminant analysis effect size (LEfSe) detected taxa differentially abundant according the ecosystems. LEfSe showed the habitat specificity of genera that were specifically more abundant in either the lake or the lagoon (Fig. 4b). This observation was consistent with previous reports showing habitat specificity for microbial taxa (Jeanbille et al., 2016a; Jeanbille et al., 2016b; Xi et al., 2014). Similarly, LEfSe revealed the presence of specific genera (*Sulfurimonas*, *Acidovorax*, *Pelobacter*, *Sediminibacter*, *Sinobacteraceae*, *Desulfobulbus*, and *Sulfurovum*) in the contaminated stations (Fig. 4c). These observations were further supported by clustering and SIMPER analyses. Clustering analysis, based on Pearson correlation, showed that the lake stations had bacterial communities with different composition to those of the lagoon stations (Fig. 5), which was corroborated by SIMPER analysis (50%). The bacterial communities from the lake stations ST5 and ST8 showed more similar composition (SIMPER: 93%) than those of the lagoon stations ST12 and ST17 (SIMPER: 71%). Consistent with LEfSe, the less contaminated station ST17 exhibited specific dominant OTUs affiliated to *Rhodobacter*, *Planctomycetaceae*, *Rhizobium*, *Rhodopirellula*, *Blastopirellula*, *Nitrosococcus* and *Sorangium* genera (Fig. 5), which related members are typically detected in marine ecosystems (Ben Said et al., 2008; Brinkhoff et al., 2012; Hernandez-Raquet et al., 2006; Jasmin et al., 2017; Klotz et al., 2006). Regarding the contaminated stations, the clustering analysis allowed the identification of specific OTUs for each station. The most PAH-contaminated station ST12 was dominated by specific OTUs belonging to *Sediminibacter* and *Loktanella* genera and other genera affiliated to *Rhodobacteraceae*, *Flavobacteriaceae* and *Saprospiraceae* families (Fig. 5), which related members have been detected in marine sediments after oil spill (Acosta-Gonzalez et al., 2015; Galazka et al., 2018; Jimenez et al., 2007). The lake stations ST5 and ST8 were dominated by specific OTUs related to *Thiobacillus*, *Nitrospira* and *Acidovorax* genera, *Escherichia-Shigella* group, and other genera affiliated to *Gemmatimonadaceae*, *Nitrospiraceae*,

*Caldilineaceae* and *Desulfarculaceae* families (Fig. 5), which was consistent with LEfSe (Fig. 4). Related members of these taxa have been found in sediments contaminated by hydrocarbons (Jochum et al., 2018; Patel et al., 2015; Reyes-Sosa et al., 2018) and pesticides (Douglass et al., 2015; Jiang et al., 2018).

Although LEfSe revealed that the abundance of some taxa were different between the lake and lagoon ecosystems (Fig. 4), the clustering analysis showed that the four stations shared dominant OTUs affiliated to *Acidithiobacillus*, *Desulfobulbus*, *Sinobacteraceae*, *Robiginitalea*, *Anaerolineaceae* and *Nannocystineae* (Fig. 5). Members of the *Acidithiobacillus* genera have been detected in metals contaminated ecosystems (Giloteaux et al., 2010), particularly in sediments (Fonti et al., 2013). It is not surprising to detect metal-resistant bacteria in the Ichkeul Lake / Bizerte Lagoon hydrological system since high levels of metal contents have been detected in the sediments (Ben Said et al., 2009; Zaaboub et al., 2015). The detection of members related to *Desulfobulbus*, *Robiginitalea*, *Sinobacteraceae*, *Anaerolineaceae* and *Nannocystineae* encompassing marine species, further support the ecological continuum between the Ichkeul Lake and Bizerte Lagoon. Among them some have been found in contaminated sediments bearing the capacity to degrade hydrocarbons (Li et al., 2017; Yan et al., 2018), and to resist and transform metals (Meng et al., 2019; Wang et al., 2018).

### **3.4. Influence of contamination on bacterial assemblages**

In order to further decipher the associations between bacterial genera and environmental parameters clustered correlation (ClusCor) analysis was performed at the genus level derived from the OTU abundance table (Fig. 6). First, sequences related to 11 genera including *Sulfurovum*, *Pseudomonas*, *Leptospirillum*, *Sulfurimonas*, *Acidovorax*, and *Nitrospira* genera, and genera related to *Anaerolineaceae*, *Desulfarculaceae*, *Caldilineaceae*, *Gemmatimonadaceae*, and *Nitrospiraceae* compose the Group1a. The abundance of the Group1a genera showed strong ( $0.6 > \text{Pearson correlation} < 0.8$ ) to very strong (Pearson

correlation > 0.8) positive correlations with the concentration of the different pesticides  
 (PCBs, Lindane, and other OCPs), and strong (-0.6 > Pearson correlation > -0.8) to very strong  
 (Pearson correlation < -0.8) negative correlation (exclusion) with PAHs concentrations. This  
 Group1a was closely related with the Group1b including *Desulfobulbus* and *Thiobacillus*  
 genera, and genera related to *Nannocystineae*, *Sinobacteraceae*, and *Escherichia-Shigella*,  
 which abundances showed strong (0.6 > Pearson correlation < 0.8) to very strong (Pearson  
 correlation > 0.8) positive correlations with the concentrations of pesticide compounds and  
 strong negative correlation (-0.6 > Pearson correlation > -0.8, i.e. exclusion) with the other  
 pollutants. Consistently, LEfSe identified all these genera (Group1a and Group1b)  
 specifically abundant for either the lake or the contamination (Fig. 4). Several *Pseudomonas*  
 and *Acidovorax* species have been isolated for their ability to degrade pesticides and other  
 organic compounds in diverse habitats (Marican and Duran-Lara, 2018; Song et al., 2000).  
 Interestingly, the presence of the sulfur-oxidizing bacteria related to the *Sulfurovum* and  
*Sulfurimonas* genera suggested the potential anaerobic degradation of pesticides since these  
 genera have been described for their capacity to perform reductive dechlorination of PCB in  
 marine sediments (Matturro et al., 2016b), often associated with sulfate-reducing bacteria  
 (SRB). Also, both genera, *Sulfurovum* and *Sulfurimonas*, have been shown to be able to grow  
 under microaerophilic conditions using oxygen or nitrate as electron acceptor (Mitchell et al.,  
 2014; Xu et al., 2017). It is likely that these genera in association with the SRB  
*Desulfarculaceae* related genera and *Desulfobulbus* in the Group1 assemblage might play a  
 role in the sulfur cycle. They probably use the hydrogen sulfur (H<sub>2</sub>S) produced by SRB and  
 oxygen for their growth playing a role in the degradation of pesticides. The genera related to  
 the *Nitrospiraceae* have been found dominant in contaminated aquatic ecosystems:  
*Nitrospira* dominant in organic matter contaminated waters (Huang et al., 2017; Wang et al.,  
 2016), while *Leptospirillum* in the presence of metals and metalloids (Wang et al., 2013).

Also, the presence of *Nitrospira* together with *Thiobacillus*, *Gemmatimonadaceae*, and the genera of the phylum Chloroflexi (*Anaerolineaceae* and *Caldilineaceae*), denitrifying genera, suggested a contamination from wastewater treatment systems as previously reported (Aanderud et al., 2018; Zhang et al., 2017), and further supported by the presence of *Escherichia-Shigella* and *Sinobacteraceae* related genera (Zhang et al., 2018). Related members of the *Nannocystineae* genera have been described harbouring the capacity to degrade complex macromolecules (Garcia and Müller, 2013). They have been detected in polluted sediments (Acosta-González et al., 2013), but their potential for pollutant degradation still to be demonstrated.

The Group2 included the genera related to *Flavobacteriaceae*, *Flammeovirgaceae*, *Saprospiraceae*, *Rhodobacteraceae* families, and the genera *Actibacter*, *Sediminibacter*, *Robiginitalea*, *Loktanella*, *Pelobacter*, *Desulfuromonas*. The abundances of Group2 genera showed very strong positive correlation with the concentrations of PAHs and organic-SN compounds (Pearson correlation >0.8), and strong (-0.6 > Pearson correlation > -0.8) to very strong (Pearson correlation < -0.8) negative correlations (exclusion) with the concentrations of lindane and other OCPs. These genera, which abundance was correlated to either the lagoon or the contamination level by LefSe (Fig. 4), have been found in hydrocarbon contaminated marine sediments (Acosta-Gonzalez et al., 2015; Chen See et al., 2018; Galazka et al., 2018) and oil production facilities (Vigneron et al., 2016; Zapata-Peñasco et al., 2013). Particularly interesting, the presence of members related to the *Desulfuromonadaceae* family (*Pelobacter* and *Desulfuromonas*) in this Group2 assemblage that have been described in presence of pesticide and for their capacity of reductive dechlorination (Greene, 2014; Sercu et al., 2013). Such observation suggested that these genera might play a key role to cope with the multi-contamination in lagoon sediments.

Behind these two major groups, the ClusCor analyses identified bacterial assemblages (Group3) including genera, which abundances showed mainly strong ( $-0.6 < \text{Pearson correlation} < -0.8$ ) to very strong ( $\text{Pearson correlation} < -0.8$ ) negative correlations (exclusion) with pollutant contents (Fig. 6). The Group3a included genera which abundance showed exclusion with PCBs and PAHs. The abundance of genera from Group3b showed exclusion with the content of all pesticides except heptachlor and strong positive correlation ( $0.6 < \text{Pearson correlation} < 0.8$ ) with PAHs content (Fig.6). The Group3c correspond to genera which abundance showed exclusion with PCBs content, while the abundance of genera from Group3d showed exclusion with PCBs and lindane contents (Fig. 6). Noteworthy, most of the genera included in Group3c (*Candidatus-Thiobios*, *Rhizobium*, *Desulfocapsa*, *Planctomycetaceae*, *Rhodopirellula*, and *Blastopirellula*) showed exclusion with pollutants, which was consistent with LEfSe highlighting that the abundance of these genera was correlated with low contamination level (Fig. 4c). It is likely that these genera were sensitive to the presence of pollutants suggesting that their presence might report on ecosystem quality. However, further investigations are requested to understand their role in the ecosystem functioning.

The co-occurrence analyses showed specific bacterial assemblages linked with contaminants revealing possible metabolic capacities of bacterial genera. Our results are consistent with previous reports that revealed specific populations linked with pollutants in marine ecosystems (Duran et al., 2015a), in freshwater lakes (Warnecke et al., 2005), and in soils (Llado et al., 2015). Such information provides new insights useful to guide future research to answer important questions in the field of microbial ecology such as the role of specific populations in the ecosystem functioning and pollutants degradation, and the role of environmental parameters in driving bacterial assemblages.

#### **4. Conclusions**

The determination of bacterial composition in sediment of the Ichkeul Lake / Bizerte Lagoon hydrological system showed that the lake bacterial communities were distinct from those of the lagoon. Although the difference could be attributed to physical-chemical parameters, the salinity and the presence of pollutants playing an important role in structuring the bacterial communities, the study revealed that the lake and the lagoon shared some bacterial genera. Such result suggested a lake-lagoon continuum due to the hydrology of the system exchanging waters according to the season resulting in bacterial community coalescence. It is likely that such bacterial genera ensure universal biogeochemical functions and, therefore constitute "generalists" bacteria essential for the basic ecosystem functioning. The study also revealed "specialists" bacteria either for the habitat (lake or lagoon) or for the pollutants. Bacterial "specialists" able to cope with the presence of pesticides in the lake and bacterial "specialists" for PAHs and organic forms of Sn in the Lagoon. These findings represent valuable information, particularly the bacterial "specialists", which represent yet an unexplored metabolic potential for the development of bioremediation processes to mitigate the effect of pollutants in aquatic ecosystems. The study also revealed bacterial genera, which abundance showed exclusion (negative correlation), with the pollutants concentrations. They represent potential bacterial biomarkers reporting on ecosystem quality, which will be useful to develop bacterial tools for monitoring the effect of countermeasures taken for mitigating the pollution.

## **Acknowledgements**

The authors would like to thank Mme Nabiha M'barek (National Agency of Environment Protection Tunisia) and Mr. Taher Kaabi (Commissariat Regional Commission for Agricultural Development of Bizerte) for their logistic contribution. We acknowledge the Regional Platform for Environmental Microbiology PREMICE supported by the Aquitaine Regional Government Council and the urban community of Pau-Pyrénées.

502

503 **Funding information**

504 This study was funded by the European programme ERANET-MED through the  
505 AQUASALT (NMED-0003-01) project.

506

507 **Compliance with ethical standards**

508 **Conflict of interest**

509 The authors declare that they have no conflict of interest.

510 **Ethical statement**

511 This article does not contain any studies with human participants or animals performed by any  
512 of the authors.

513

## References

- Aanderud, Z.T., Saurey, S., Ball, B.A., Wall, D.H., Barrett, J.E., Muscarella, M.E., Griffin, N.A., Virginia, R.A., Adams, B.J., 2018. Stoichiometric Shifts in Soil C:N:P Promote Bacterial Taxa Dominance, Maintain Biodiversity, and Deconstruct Community Assemblages. *Front Microbiol* 9, 1401.
- Acosta-Gonzalez, A., Martirani-von Abercron, S.M., Rossello-Mora, R., Wittich, R.M., Marques, S., 2015. The effect of oil spills on the bacterial diversity and catabolic function in coastal sediments: a case study on the Prestige oil spill. *Environ Sci Pollut R* 22, 15200-15214.
- Acosta-González, A., Rosselló-Móra, R., Marqués, S., 2013. Characterization of the anaerobic microbial community in oil-polluted subtidal sediments: aromatic biodegradation potential after the Prestige oil spill. *Environmental Microbiology* 15, 77-92.
- Andrews, D.M., Lin, H., Zhu, Q., Jin, L.X., Brantley, S.L., 2011. Hot Spots and Hot Moments of Dissolved Organic Carbon Export and Soil Organic Carbon Storage in the Shale Hills Catchment. *Vadose Zone J* 10, 943-954.
- Bakke, T., Källqvist, T., Ruus, A., Breedveld, G.D., Hylland, K., 2010. Development of sediment quality criteria in Norway. *Journal of Soils and Sediments* 10, 172-178.
- Barberan, A., Bates, S.T., Casamayor, E.O., Fierer, N., 2012. Using network analysis to explore co-occurrence patterns in soil microbial communities. *ISME J* 6, 343-351.
- Behera, P., Mahapatra, S., Mohapatra, M., Kim, J.Y., Adhya, T.K., Raina, V., Suar, M., Pattnaik, A.K., Rastogi, G., 2017. Salinity and macrophyte drive the biogeography of the sedimentary bacterial communities in a brackish water tropical coastal lagoon. *Science of the Total Environment* 595, 472-485.
- Ben Said, O., Goñi-Urriza, M., El Bour, M., Aissa, P., Duran, R., 2009. Bacterial Community Structure of Sediments of the Bizerte Lagoon (Tunisia), a Southern Mediterranean Coastal Anthropized Lagoon. *Microb Ecol* 59, 445-456.
- Ben Said, O., Goni-Urriza, M.S., El Bour, M., Dellali, M., Aissa, P., Duran, R., 2008. Characterization of aerobic polycyclic aromatic hydrocarbon-degrading bacteria from Bizerte lagoon sediments, Tunisia. *J Appl Microbiol* 104, 987-997.
- Ben Said, O., Louati, H., Soltani, A., Preud'homme, H., Cravo-Laureau, C., Got, P., Pringault, O., Aissa, P., Duran, R., 2015. Changes of benthic bacteria and meiofauna assemblages during bio-treatments of anthracene-contaminated sediments from Bizerta lagoon (Tunisia). *Environ Sci Pollut R* 22, 15319-15331.
- Ben Salem, F., Ben Said, O., Aissa, P., Mahmoudi, E., Monperrus, M., Grunberger, O., Duran, R., 2016a. Pesticides in Ichkeul Lake-Bizerta Lagoon Watershed in Tunisia: use, occurrence, and effects on bacteria and free-living marine nematodes. *Environ Sci Pollut R* 23, 36-48.
- Ben Salem, F., Ben Said, O., Duran, R., Monperrus, M., 2016b. Validation of an Adapted QuEChERS Method for the Simultaneous Analysis of Polycyclic Aromatic Hydrocarbons, Polychlorinated Biphenyls and Organochlorine Pesticides in Sediment by Gas Chromatography-Mass Spectrometry. *Bull Environ Contam Toxicol* 96, 678-684.
- Ben Salem, F., Ben Said, O., Mahmoudi, E., Duran, R., Monperrus, M., 2017. Distribution of organic contamination of sediments from Ichkeul Lake and Bizerte Lagoon, Tunisia. *Mar Pollut Bull* 123, 329-338.
- Berg, J., Brandt, K.K., Al-Soud, W.A., Holm, P.E., Hansen, L.H., Sørensen, S.J., Nybroe, O., 2012. Selection for Cu-tolerant bacterial communities with altered composition, but unaltered richness, via long-term Cu exposure. *Applied and Environmental Microbiology* 78, 7438-7446.



563 Bodoque, J.M., Ladera, J., Yela, J.L., Alonso-Azcárate, J., Brito, D., Antigüedad, I., Duran,  
 564 R., Attard, E., Lauga, B., Sánchez-Pérez, J.M., 2017. Recovering hydromorphological  
 565 functionality to improve natural purification capacity of a highly human-modified wetland.  
 566 *Ecological Engineering* 103, 332-343.  
 567 Bordenave, S., Fourcans, A., Blanchard, S., Goni, M.S., Caumette, P., Duran, R., 2004a.  
 568 Structure and functional analyses of bacterial communities changes in microbial mats  
 569 following petroleum exposure. *Ophelia* 58, 195-203.  
 570 Bordenave, S., Goni-Urriza, M., Vilette, C., Blanchard, S., Caumette, P., Duran, R., 2008.  
 571 Diversity of ring-hydroxylating dioxygenases in pristine and oil contaminated microbial mats  
 572 at genomic and transcriptomic levels. *Environ Microbiol* 10, 3201-3211.  
 573 Bordenave, S., Jézéquel, R., Fourcans, A., Budzinski, H., Merlin, F.X., Fourel, T., Goñi-  
 574 Urriza, M., Guyoneaud, R., Grimaud, R., Caumette, P., Duran, R., 2004b. Degradation of  
 575 the "Erika" oil. *Aquat Living Resour* 17, 261-267.  
 576 Borin, S., Brusetti, L., Daffonchio, D., Delaney, E., Baldi, F., 2009. Biodiversity of  
 577 prokaryotic communities in sediments of different sub-basins of the Venice lagoon. *Research*  
 578 *in Microbiology* 160, 307-314.  
 579 Botta, F., Lavison, G., Couturier, G., Alliot, F., Moreau-Guigon, E., Fauchon, N., Guery, B.,  
 580 Chevreuil, M., Blanchoud, H., 2009. Transfer of glyphosate and its degradate AMPA to  
 581 surface waters through urban sewerage systems. *Chemosphere* 77, 133-139.  
 582 Brinkhoff, T., Fischer, D., Vollmers, J., Voget, S., Beardsley, C., Thole, S., Musmann, M.,  
 583 Kunze, B., Wagner-Dobler, I., Daniel, R., Simon, M., 2012. Biogeography and phylogenetic  
 584 diversity of a cluster of exclusively marine myxobacteria. *ISME J* 6, 1260-1272.  
 585 Brito, E.M., Duran, R., Guyoneaud, R., Goni-Urriza, M., Garcia de Oteyza, T., Crapez, M.A.,  
 586 Aleluia, I., Wasserman, J.C., 2009. A case study of in situ oil contamination in a mangrove  
 587 swamp (Rio De Janeiro, Brazil). *Mar Pollut Bull* 58, 418-423.  
 588 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K.,  
 589 Fierer, N., Pena, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D.,  
 590 Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder,  
 591 J., Sevinsky, J.R., Tumbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J.,  
 592 Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data. *Nat*  
 593 *Methods* 7, 335-336.  
 594 Casagrande, C., Boudouresque, C.F., 2010. A first quantification of the overall biomass, from  
 595 phytoplankton to birds, of a Mediterranean brackish lagoon: Revisiting the ecosystem of Lake  
 596 Ichkeul (Tunisia). *Hydrobiologia* 637, 73-85.  
 597 Chang, S.M., Doong, R.A., 2006. Concentration and fate of persistent organochlorine  
 598 pesticides in estuarine sediments using headspace solid-phase microextraction. *Chemosphere*  
 599 62, 1869-1878.  
 600 Chen See, J.R., Ulrich, N., Nwanosike, H., McLimans, C.J., Tokarev, V., Wright, J.R.,  
 601 Campa, M.F., Grant, C.J., Hazen, T.C., Niles, J.M., Ressler, D., Lamendella, R., 2018.  
 602 Bacterial Biomarkers of Marcellus Shale Activity in Pennsylvania. *Front Microbiol* 9, 1697.  
 603 Chronopoulou, P.M., Fahy, A., Coulon, F., Paise, S., Goni-Urriza, M., Peperzak, L., Alvarez,  
 604 L.A., McKew, B.A., Lawson, T., Timmis, K.N., Duran, R., Underwood, G.J.C., McGenity,  
 605 T.J., 2013. Impact of a simulated oil spill on benthic phototrophs and nitrogen-fixing bacteria  
 606 in mudflat mesocosms. *Environ Microbiol* 15, 242-252.  
 607 Cravo-Laureau, C., Cagnon, C., Lauga, B., Duran, R., 2017a. Preface. *Microbial*  
 608 *Ecotoxicology*, v-vi.  
 609 Cravo-Laureau, C., Duran, R., 2014. Marine coastal sediments microbial hydrocarbon  
 610 degradation processes: contribution of experimental ecology in the omics'era. *Front Microbiol*  
 611 5, 39.

612 Cravo-Laureau, C., Hernandez-Raquet, G., Vitte, I., Jezequel, R., Bellet, V., Godon, J.J.,  
613 Caumette, P., Balaguer, P., Duran, R., 2011. Role of environmental fluctuations and microbial  
614 diversity in degradation of hydrocarbons in contaminated sludge. *Res Microbiol* 162, 888-  
615 895.

616 Cravo-Laureau, C., Lauga, B., Cagnon, C., Duran, R., 2017b. Microbial responses to  
617 pollution- ecotoxicology: Introducing the different biological levels, *Microbial*  
618 *Ecotoxicology*, pp. 45-62.

619 Daims, H., 2014. The family nitrospiraceae, *The Prokaryotes: Other Major Lineages of*  
620 *Bacteria and The Archaea*, pp. 733-749.

621 Dell'Anno, A., Beolchini, F., Rocchetti, L., Luna, G.M., Danovaro, R., 2012. High bacterial  
622 biodiversity increases degradation performance of hydrocarbons during bioremediation of  
623 contaminated harbor marine sediments. *Environmental Pollution* 167, 85-92.

624 DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T.,  
625 Dalevi, D., Hu, P., Andersen, G.L., 2006. Greengenes, a chimera-checked 16S rRNA gene  
626 database and workbench compatible with ARB. *Applied and Environmental Microbiology* 72,  
627 5069-5072.

628 Ding, Q., Huang, X., Hu, H., Hong, M., Zhang, D., Wang, K., 2017. Impact of pyrene and  
629 cadmium co-contamination on prokaryotic community in coastal sediment microcosms.  
630 *Chemosphere* 188, 320-328.

631 Dollhopf, S.L., Hashsham, S.A., Tiedje, J.M., 2001. Interpreting 16S rDNA T-RFLP data:  
632 Application of self-organizing maps and principal component analysis to describe community  
633 dynamics and convergence. *Microbial Ecology* 42, 495-505.

634 Dominique, Y., Muresan, B., Duran, R., Richard, S., Boudou, A., 2007. Simulation of the  
635 chemical fate and bioavailability of liquid elemental mercury drops from gold mining in  
636 Amazonian freshwater systems. *Environmental Science and Technology* 41, 7322-7329.

637 Doong, R.A., Lee, S.H., Lee, C.C., Sun, Y.C., Wu, S.C., 2008. Characterization and  
638 composition of heavy metals and persistent organic pollutants in water and estuarine  
639 sediments from Gao-ping River, Taiwan. *Mar Pollut Bull* 57, 846-857.

640 Douglass, J.F., Radosevich, M., Tuovinen, O.H., 2015. Molecular analysis of atrazine-  
641 degrading bacteria and catabolic genes in the water column and sediment of a created wetland  
642 in an agricultural/urban watershed. *Ecol Eng* 83, 405-412.

643 Dowd, S.E., Callaway, T.R., Wolcott, R.D., Sun, Y., McKeenhan, T., Hagevoort, R.G.,  
644 Edrington, T.S., 2008. Evaluation of the bacterial diversity in the feces of cattle using 16S  
645 rDNA bacterial tag-encoded FLX amplicon pyrosequencing (bTEFAP). *BMC Microbiol* 8,  
646 125.

647 Duran, R., Bielen, A., Paradzik, T., Gassie, C., Pustijanac, E., Cagnon, C., Hamer, B.,  
648 Vujaklija, D., 2015a. Exploring Actinobacteria assemblages in coastal marine sediments  
649 under contrasted Human influences in the West Istria Sea, Croatia. *Environ Sci Pollut R* 22,  
650 15215-15229.

651 Duran, R., Bonin, P., Jezequel, R., Dubosc, K., Gassie, C., Terrisse, F., Abella, J., Cagnon, C.,  
652 Milton, C., Michotey, V., Gilbert, F., Cuny, P., Cravo-Laureau, C., 2015b. Effect of physical  
653 sediments reworking on hydrocarbon degradation and bacterial community structure in  
654 marine coastal sediments. *Environ Sci Pollut R* 22, 15248-15259.

655 Duran, R., Cravo-Laureau, C., 2016. Role of environmental factors and microorganisms in  
656 determining the fate of polycyclic aromatic hydrocarbons in the marine environment. *FEMS*  
657 *Microbiol Rev* 40, 814-830.

658 Duran, R., Cuny, P., Bonin, P., Cravo-Laureau, C., 2015c. Microbial ecology of hydrocarbon-  
659 polluted coastal sediments. *Environ Sci Pollut R* 22, 15195-15199.

660 Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST.  
661 *Bioinformatics* 26, 2460-2461.

Fagervold, S.K., May, H.D., Sowers, K.R., 2007. Microbial reductive dechlorination of Aroclor 1260 in Baltimore Harbor sediment microcosms is catalyzed by three phylotypes within the Phylum chloroflexi. *Applied and Environmental Microbiology* 73, 3009-3018.

Fan, L., Barry, K., Hu, G., Meng, S., Song, C., Qiu, L., Zheng, Y., Wu, W., Qu, J., Chen, J., Xu, P., 2016. Characterizing bacterial communities in tilapia pond surface sediment and their responses to pond differences and temporal variations. *World Journal of Microbiology and Biotechnology* 33, 1.

Fernanda Sánchez-Soto Jiménez, M., Cerqueda-García, D., Montero-Muñoz, J.L., Leopoldina Aguirre-Macedo, M., García-Maldonado, J.Q., 2018. Assessment of the bacterial community structure in shallow and deep sediments of the Perdido Fold Belt region in the Gulf of Mexico. *PeerJ* 2018.

Fonti, V., Dell'Anno, A., Beolchini, F., 2013. Influence of biogeochemical interactions on metal bioleaching performance in contaminated marine sediment. *Water Res* 47, 5139-5152.

Fourcans, A., Ranchou-Peyruse, A., Caumette, P., Duran, R., 2008. Molecular analysis of the spatio-temporal distribution of sulfate-reducing bacteria (SRB) in Camargue (France) hypersaline microbial mat. *Microb Ecol* 56, 90-100.

Fuhrman, J.A., Steele, J.A., 2008. Community structure of marine bacterioplankton: patterns, networks, and relationships to function. *Aquat Microb Ecol* 53, 69-81.

Galazka, A., Grzadziel, J., Galazka, R., Ukalska-Jaruga, A., Strzelecka, J., Smreczak, B., 2018. Genetic and Functional Diversity of Bacterial Microbiome in Soils With Long Term Impacts of Petroleum Hydrocarbons. *Front Microbiol* 9, 1923.

Garcia, R., Müller, R., 2013. The family nannocystaceae, *The Prokaryotes: Deltaproteobacteria and Epsilonproteobacteria*, pp. 213-229.

Gharbi-Bouraoui, S., Gnassia-Barelli, M., Roméo, M., Dellali, M., Aïssa, P., 2008. Field study of metal concentrations and biomarker responses in the neogastropod, *Murex trunculus*, from Bizerta Lagoon (Tunisia). *Aquatic Living Resources* 21, 213-220.

Giloteaux, L., Goni-Urriza, M., Duran, R., 2010. Nested PCR and new primers for analysis of sulfate-reducing bacteria in low-cell-biomass environments. *Appl Environ Microbiol* 76, 2856-2865.

Greene, A.C., 2014. The family desulfuromonadaceae, *The Prokaryotes: Deltaproteobacteria and Epsilonproteobacteria*, pp. 143-155.

Guermouche M'rassi, A., Bensalah, F., Gury, J., Duran, R., 2015. Isolation and characterization of different bacterial strains for bioremediation of n-alkanes and polycyclic aromatic hydrocarbons. *Environ Sci Pollut R* 22, 15332-15346.

Haller, L., Tonolla, M., Zopfi, J., Peduzzi, R., Wildi, W., Pote, J., 2011. Composition of bacterial and archaeal communities in freshwater sediments with different contamination levels (Lake Geneva, Switzerland). *Water Res* 45, 1213-1228.

Hernandez-Raquet, G., Budzinski, H., Caumette, P., Dabert, P., Le Menach, K., Muyzer, G., Duran, R., 2006. Molecular diversity studies of bacterial communities of oil polluted microbial mats from the Etang de Berre (France). *FEMS Microbiol Ecol* 58, 550-562.

Huang, W., Chen, X., Jiang, X., Zheng, B.H., 2017. Characterization of sediment bacterial communities in plain lakes with different trophic statuses. *MicrobiologyOpen* 6.

Hug, L.A., Castelle, C.J., Wrighton, K.C., Thomas, B.C., Sharon, I., Frischkorn, K.R., Williams, K.H., Tringe, S.G., Banfield, J.F., 2013. Community genomic analyses constrain the distribution of metabolic traits across the Chloroflexi phylum and indicate roles in sediment carbon cycling. *Microbiome* 1.

Hunter, E.M., Mills, H.J., Kostka, J.E., 2006. Microbial community diversity associated with carbon and nitrogen cycling in permeable shelf sediments. *Applied and Environmental Microbiology* 72, 5689-5701.

711 Ikenaga, M., Guevara, R., Dean, A.L., Pisani, C., Boyer, J.N., 2010. Changes in Community  
 712 Structure of Sediment Bacteria Along the Florida Coastal Everglades Marsh-Mangrove-  
 713 Seagrass Salinity Gradient. *Microb Ecol* 59, 284-295.  
 714 Jacquiod, S., Cyriaque, V., Riber, L., Al-soud, W.A., Gillan, D.C., Wattiez, R., Sørensen, S.J.,  
 715 2018a. Long-term industrial metal contamination unexpectedly shaped diversity and activity  
 716 response of sediment microbiome. *Journal of Hazardous Materials* 344, 299-307.  
 717 Jacquiod, S., Nunes, I., Brejnrod, A., Hansen, M.A., Holm, P.E., Johansen, A., Brandt, K.K.,  
 718 Priemé, A., Sørensen, S.J., 2018b. Long-term soil metal exposure impaired temporal variation  
 719 in microbial metatranscriptomes and enriched active phages. *Microbiome* 6.  
 720 Jasmin, C., Anas, A., Tharakan, B., Jaleel, A., Puthiyaveetil, V., Narayanane, S., Lincy, J.,  
 721 Nair, S., 2017. Diversity of sediment-associated Planctomycetes in the Arabian Sea oxygen  
 722 minimum zone. *J Basic Microbiol* 57, 1010-1017.  
 723 Jeanbille, M., Gury, J., Duran, R., Tronczynski, J., Agogue, H., Ben Said, O., Ghiglione, J.F.,  
 724 Auguet, J.C., 2016a. Response of Core Microbial Consortia to Chronic Hydrocarbon  
 725 Contaminations in Coastal Sediment Habitats. *Front Microbiol* 7, 1637.  
 726 Jeanbille, M., Gury, J., Duran, R., Tronczynski, J., Ghiglione, J.F., Agogue, H., Said, O.B.,  
 727 Taib, N., Debroas, D., Garnier, C., Auguet, J.C., 2016b. Chronic Polyaromatic Hydrocarbon  
 728 (PAH) Contamination Is a Marginal Driver for Community Diversity and Prokaryotic  
 729 Predicted Functioning in Coastal Sediments. *Front Microbiol* 7, 1303.  
 730 Jiang, L., Luo, C., Zhang, D., Song, M., Sun, Y., Zhang, G., 2018. Biphenyl-Metabolizing  
 731 Microbial Community and a Functional Operon Revealed in E-Waste-Contaminated Soil.  
 732 *Environ Sci Technol* 52, 8558-8567.  
 733 Jimenez, N., Vinas, M., Bayona, J.M., Albaiges, J., Solanas, A.M., 2007. The Prestige oil  
 734 spill: bacterial community dynamics during a field biostimulation assay. *Appl Microbiol*  
 735 *Biotechnol* 77, 935-945.  
 736 Jochum, L.M., Schreiber, L., Marshall, I.P.G., Jorgensen, B.B., Schramm, A., Kjeldsen, K.U.,  
 737 2018. Single-Cell Genomics Reveals a Diverse Metabolic Potential of Uncultivated  
 738 Desulfatiglans-Related Deltaproteobacteria Widely Distributed in Marine Sediment. *Front*  
 739 *Microbiol* 9.  
 740 Johnston, E.L., Roberts, D.A., 2009. Contaminants reduce the richness and evenness of  
 741 marine communities: a review and meta-analysis. *Environ Pollut* 157, 1745-1752.  
 742 Junier, P., Molina, V., Dorador, C., Hadas, O., Kim, O.S., Junier, T., Witzel, K.P., Imhoff,  
 743 J.F., 2010. Phylogenetic and functional marker genes to study ammonia-oxidizing  
 744 microorganisms (AOM) in the environment. *Applied Microbiology and Biotechnology* 85,  
 745 425-440.  
 746 Jurgburg, S.D., Nunes, I., Brejnrod, A., Jacquiod, S., Priemé, A., Sørensen, S.J., Van Elsas,  
 747 J.D., Salles, J.F., 2017. Legacy Effects on the Recovery of Soil Bacterial Communities from  
 748 Extreme Temperature Perturbation. *Frontiers in Microbiology* 8.  
 749 Kjellerup, B.V., Naff, C., Edwards, S.J., Ghosh, U., Baker, J.E., Sowers, K.R., 2014. Effects  
 750 of activated carbon on reductive dechlorination of PCBs by organohalide respiring bacteria  
 751 indigenous to sediments. *Water Res* 52, 1-10.  
 752 Klotz, M.G., Arp, D.J., Chain, P.S.G., El-Sheikh, A.F., Hauser, L.J., Hommes, N.G., Larimer,  
 753 F.W., Malfatti, S.A., Norton, J.M., Poret-Peterson, A.T., Vergez, L.M., Ward, B.B., 2006.  
 754 Complete Genome Sequence of the Marine, Chemolithoautotrophic, Ammonia-Oxidizing  
 755 Bacterium *Nitrosococcus oceanus* ATCC 19707. *Appl Environ Microbiol* 72, 6299-6315.  
 756 Li, J., Zhang, D., Song, M., Jiang, L., Wang, Y., Luo, C., Zhang, G., 2017. Novel bacteria  
 757 capable of degrading phenanthrene in activated sludge revealed by stable-isotope probing  
 758 coupled with high-throughput sequencing. *Biodegradation* 28, 423-436.

759 Lipsewers, Y.A., Hopmans, E.C., Meysman, F.J.R., Sinninghe Damsté, J.S., Villanueva, L.,  
 760 2016. Abundance and Diversity of Denitrifying and Anammox Bacteria in Seasonally  
 761 Hypoxic and Sulfidic Sediments of the Saline Lake Grevelingen. *Frontiers in Microbiology* 7.  
 762 Liu, S., Ren, H., Shen, L., Lou, L., Tian, G., Zheng, P., Hu, B., 2015. pH levels drive bacterial  
 763 community structure in sediments of the Qiantang River as determined by 454  
 764 pyrosequencing. *Frontiers in microbiology* 6, 285-285.  
 765 Llado, S., Covino, S., Solanas, A.M., Petruccioli, M., D'Annibale, A., Vinas, M., 2015.  
 766 Pyrosequencing reveals the effect of mobilizing agents and lignocellulosic substrate  
 767 amendment on microbial community composition in a real industrial PAH-polluted soil. *J*  
 768 *Hazard Mater* 283, 35-43.  
 769 Long, E.R., Field, L.J., MacDonald, D.D., 1998. Predicting toxicity in marine sediments with  
 770 numerical sediment quality guidelines. *Environmental Toxicology and Chemistry* 17, 714-  
 771 727.  
 772 Louati, H., Ben Said, O., Got, P., Soltani, A., Mahmoudi, E., Cravo-Laureau, C., Duran, R.,  
 773 Aissa, P., Pringault, O., 2013a. Microbial community responses to bioremediation treatments  
 774 for the mitigation of low-dose anthracene in marine coastal sediments of Bizerte lagoon  
 775 (Tunisia). *Environ Sci Pollut R* 20, 300-310.  
 776 Louati, H., Said, O.B., Soltani, A., Got, P., Mahmoudi, E., Cravo-Laureau, C., Duran, R.,  
 777 Aissa, P., Pringault, O., 2013b. The roles of biological interactions and pollutant  
 778 contamination in shaping microbial benthic community structure. *Chemosphere* 93, 2535-  
 779 2546.  
 780 Mahmoudi, N., Robeson, M.S., II, Castro, H.F., Fortney, J.L., Techtmann, S.M., Joyner, D.C.,  
 781 Paradis, C.J., Pfiffner, S.M., Hazen, T.C., 2015. Microbial community composition and  
 782 diversity in Caspian Sea sediments. *FEMS Microbiology Ecology* 91.  
 783 Mansour, I., Heppell, C.M., Ryo, M., Rillig, M.C., 2018. Application of the microbial  
 784 community coalescence concept to riverine networks. *Biological Reviews* 93, 1832-1845.  
 785 Marchesi, J.R., Sato, T., Weightman, A.J., Martin, T.A., Fry, J.C., Hiom, S.J., Wade, W.G.,  
 786 1998. Design and evaluation of useful bacterium-specific PCR primers that amplify genes  
 787 coding for bacterial 16S rRNA. *Appl Environ Microbiol* 64, 795-799.  
 788 Marican, A., Duran-Lara, E.F., 2018. A review on pesticide removal through different  
 789 processes. *Environ Sci Pollut R* 25, 2051-2064.  
 790 Matturro, B., Presta, E., Rossetti, S., 2016a. Reductive dechlorination of tetrachloroethene in  
 791 marine sediments: Biodiversity and dehalorespiring capabilities of the indigenous microbes.  
 792 *Science of the Total Environment* 545-546, 445-452.  
 793 Matturro, B., Ubaldi, C., Rossetti, S., 2016b. Microbiome Dynamics of a Polychlorobiphenyl  
 794 (PCB) Historically Contaminated Marine Sediment under Conditions Promoting Reductive  
 795 Dechlorination. *Front Microbiol* 7, 1502.  
 796 McDonald, D., Price, M.N., Goodrich, J., Nawrocki, E.P., Desantis, T.Z., Probst, A.,  
 797 Andersen, G.L., Knight, R., Hugenholtz, P., 2012. An improved Greengenes taxonomy with  
 798 explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME Journal*  
 799 6, 610-618.  
 800 Meng, D., Li, J., Liu, T., Liu, Y., Yan, M., Hu, J., Li, X., Liu, X., Liang, Y., Liu, H., Yin, H.,  
 801 2019. Effects of redox potential on soil cadmium solubility: Insight into microbial  
 802 community. *J Environ Sci (China)* 75, 224-232.  
 803 Misson, B., Garnier, C., Lauga, B., Dang, D.H., Ghiglione, J.F., Mullot, J.U., Duran, R.,  
 804 Pringault, O., 2016. Chemical multi-contamination drives benthic prokaryotic diversity in the  
 805 anthropized Toulon Bay. *Sci Total Environ* 556, 319-329.  
 806 Mitchell, H.M., Rocha, G.A., Kaakoush, N.O., O'Rourke, J.L., Queiroz, D.M.M., 2014. The  
 807 family helicobacteraceae, *The Prokaryotes: Deltaproteobacteria and Epsilonproteobacteria*,  
 808 pp. 337-392.

809 Mwanamoki, P.M., Devarajan, N., Thevenon, F., Birane, N., de Alencastro, L.F., Grandjean,  
810 D., Mpiana, P.T., Prabakar, K., Mubedi, J.I., Kabele, C.G., Wildi, W., Pote, J., 2014. Trace  
811 metals and persistent organic pollutants in sediments from river-reservoir systems in  
812 Democratic Republic of Congo (DRC): Spatial distribution and potential ecotoxicological  
813 effects. *Chemosphere* 111, 485-492.

814 Nunes, I., Jurgburg, S., Jacquiod, S., Brejnrod, A., Falcão Salles, J., Priemé, A., Sørensen, S.J.,  
815 2018. Soil bacteria show different tolerance ranges to an unprecedented disturbance. *Biology  
816 and Fertility of Soils* 54, 189-202.

817 Obi, C.C., Adebuseye, S.A., Ugoji, E.O., Ilori, M.O., Amund, O.O., Hickey, W.J., 2016.  
818 Microbial Communities in Sediments of Lagos Lagoon, Nigeria: Elucidation of Community  
819 Structure and Potential Impacts of Contamination by Municipal and Industrial Wastes.  
820 *Frontiers in Microbiology* 7.

821 Oksanen, J., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara, R., Simpson, G.,  
822 Solymos, P., Henry, M., Stevens, H., 2013. *Vegan: Community Ecology Package*. 2013. R-  
823 package version 2.0-10. URL <http://CRAN.R-project.org/package=vegan>.

824 Paisse, S., Coulon, F., Goni-Urriza, M., Peperzak, L., McGenity, T.J., Duran, R., 2008.  
825 Structure of bacterial communities along a hydrocarbon contamination gradient in a coastal  
826 sediment. *FEMS Microbiol Ecol* 66, 295-305.

827 Paissé, S., Goñi-Urriza, M., Stadler, T., Budzinski, H., Duran, R., 2012. Ring-hydroxylating  
828 dioxygenase (RHD) expression in a microbial community during the early response to oil  
829 pollution. *FEMS Microbiology Ecology* 80, 77-86.

830 Patel, V., Munot, H., Shah, V., Shouche, Y.S., Madamwar, D., 2015. Taxonomic profiling of  
831 bacterial community structure from coastal sediment of Alang-Sosiya shipbreaking yard near  
832 Bhavnagar, India. *Mar Pollut Bull* 101, 736-745.

833 Pavloudi, C., Oulas, A., Vasileiadou, K., Sarropoulou, E., Kotoulas, G., Arvanitidis, C., 2016.  
834 Salinity is the major factor influencing the sediment bacterial communities in a Mediterranean  
835 lagoonal complex (Amvrakikos Gulf, Ionian Sea). *Marine Genomics* 28, 71-81.

836 Pérez-Domínguez, R., MacI, S., Courrat, A., Lepage, M., Borja, A., Uriarte, A., Neto, J.M.,  
837 Cabral, H., St. raykov, V., Franco, A., Alvarez, M.C., Elliott, M., 2012. Current developments  
838 on fish-based indices to assess ecological-quality status of estuaries and lagoons. *Ecological  
839 Indicators* 23, 34-45.

840 Precigou, S., Goulas, P., Duran, R., 2001. Rapid and specific identification of nitrile hydratase  
841 (NHase)-encoding genes in soil samples by polymerase chain reaction. *FEMS Microbiol Lett*  
842 204, 155-161.

843 Precigou, S., Wieser, M., Pommars, P., Goulas, P., Duran, R., 2004. *Rhodococcus*  
844 *pyridinovorans* MW3, a bacterium producing a nitrile hydratase. *Biotechnol Lett* 26, 1379-  
845 1384.

846 Pringault, O., Duran, R., Jacquet, S., Torreton, J.P., 2008. Temporal variations of microbial  
847 activity and diversity in marine tropical sediments (New Caledonia lagoon). *Microb Ecol* 55,  
848 247-258.

849 Quero, G.M., Cassin, D., Botter, M., Perini, L., Luna, G.M., 2015. Patterns of benthic  
850 bacterial diversity in coastal areas contaminated by heavy metals, polycyclic aromatic  
851 hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs). *Frontiers in Microbiology* 6.

852 Quince, C., Lanzen, A., Curtis, T.P., Davenport, R.J., Hall, N., Head, I.M., Read, L.F., Sloan,  
853 W.T., 2009. Accurate determination of microbial diversity from 454 pyrosequencing data. *Nat  
854 Methods* 6, 639-641.

855 Ramdani, M., Flower, R.J., Elkhiaiti, N., Kraïem, M.M., Fathi, A.A., Birks, H.H., Patrick,  
856 S.T., 2001. North African wetland lakes: Characterization of nine sites included in the  
857 CASSARINA Project. *Aquat Ecol* 35, 281-302.

858 Reyes-Sosa, M.B., Apodaca-Hernandez, J.E., Arena-Ortiz, M.L., 2018. Bioprospecting for  
859 microbes with potential hydrocarbon remediation activity on the northwest coast of the  
860 Yucatan Peninsula, Mexico, using DNA sequencing. *Sci Total Environ* 642, 1060-1074.

861 Rillig, M.C., Antonovics, J., Caruso, T., Lehmann, A., Powell, J.R., Veresoglou, S.D.,  
862 Verbruggen, E., 2015. Interchange of entire communities: Microbial community coalescence.  
863 *Trends in Ecology and Evolution* 30, 470-476.

864 Schultz, P., Urban, N.R., 2008. Effects of bacterial dynamics on organic matter  
865 decomposition and nutrient release from sediments: A modeling study. *Ecol Model* 210, 1-14.

866 Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., Huttenhower, C.,  
867 2011. Metagenomic biomarker discovery and explanation. *Genome Biol* 12.

868 Sercu, B., Jones, A.D., Wu, C.H., Escobar, M.H., Serlin, C.L., Knapp, T.A., Andersen, G.L.,  
869 Holden, P.A., 2013. The influence of in situ chemical oxidation on microbial community  
870 composition in groundwater contaminated with chlorinated solvents. *Microb Ecol* 65, 39-49.

871 Song, B., Palleroni, N.J., Haggblom, M.M., 2000. Isolation and characterization of diverse  
872 halobenzoate-degrading denitrifying bacteria from soils and sediments. *Appl Environ*  
873 *Microbiol* 66, 3446-3453.

874 Stauffert, M., Cravo-Laureau, C., Duran, R., 2016. Structure of hydrocarbonoclastic nitrate-  
875 reducing bacterial communities in bioturbated coastal marine sediments. *FEMS Microbiology*  
876 *Ecology* 89, 580-593.

877 Stauffert, M., Cravo-Laureau, C., Jezequel, R., Barantal, S., Cuny, P., Gilbert, F., Cagnon, C.,  
878 Militon, C., Amouroux, D., Mahdaoui, F., Bouyssiére, B., Stora, G., Merlin, F.X., Duran, R.,  
879 2013. Impact of oil on bacterial community structure in bioturbated sediments. *PLoS One* 8,  
880 e65347.

881 Stauffert, M., Duran, R., Gassie, C., Cravo-Laureau, C., 2014. Response of archaeal  
882 communities to oil spill in bioturbated mudflat sediments. *Microb Ecol* 67, 108-119.

883 Stevenson, A., Battarbee, R., 1991. Palaeoecological and documentary records of recent  
884 environmental change in Garaet El Ichkeul: a seasonally saline lake in NW Tunisia. *Biol*  
885 *Conserv* 58, 275-295.

886 Sun, M.Y., Dafforn, K.A., Johnston, E.L., Brown, M.V., 2013. Core sediment bacteria drive  
887 community response to anthropogenic contamination over multiple environmental gradients.  
888 *Environmental Microbiology* 15, 2517-2531.

889 Terrisse, F., Cravo-Laureau, C., Noel, C., Cagnon, C., Dumbrell, A.J., McGenity, T.J., Duran,  
890 R., 2017. Variation of Oxygenation Conditions on a Hydrocarbonoclastic Microbial  
891 Community Reveals *Alcanivorax* and *Cyclocasticus* Ecotypes. *Front Microbiol* 8.

892 Thiyagarajan, V., Tsoi, M.M., Zhang, W., Qian, P.Y., 2010. Temporal variation of coastal  
893 surface sediment bacterial communities along an environmental pollution gradient. *Mar*  
894 *Environ Res* 70, 56-64.

895 Trabelsi, S., Driss, M.R., 2005. Polycyclic aromatic hydrocarbons in superficial coastal  
896 sediments from Bizerte Lagoon, Tunisia. *Mar Pollut Bull* 50, 344-348.

897 Vasconcelos, R.P., Reis-Santos, P., Costa, M.J., Cabral, H.N., 2011. Connectivity between  
898 estuaries and marine environment: Integrating metrics to assess estuarine nursery function.  
899 *Ecological Indicators* 11, 1123-1133.

900 Vercraene-Eairmal, M., Lauga, B., Saint Laurent, S., Mazzella, N., Boutry, S., Simon, M.,  
901 Karama, S., Delmas, F., Duran, R., 2010. Diuron biotransformation and its effects on biofilm  
902 bacterial community structure. *Chemosphere* 81, 837-843.

903 Vigneron, A., Alsop, E.B., Chambers, B., Lomans, B.P., Head, I.M., Tsesmetzis, N., 2016.  
904 Complementary Microorganisms in Highly Corrosive Biofilms from an Offshore Oil  
905 Production Facility. *Appl Environ Microbiol* 82, 2545-2554.

Vitte, I., Duran, R., Jezequel, R., Caumette, P., Cravo-Laureau, C., 2011. Effect of oxic/anoxic switches on bacterial communities and PAH biodegradation in an oil-contaminated sludge. *Environ Sci Pollut R* 18, 1022-1032.

Wang, J., Li, Y., Wang, P., Niu, L., Zhang, W., Wang, C., 2016. Response of bacterial community compositions to different sources of pollutants in sediments of a tributary of Taihu Lake, China. *Environ Sci Pollut R* 23, 13886-13894.

Wang, J., Zhao, H.B., Zhuang, T., Qin, W.Q., Zhu, S., Qiu, G.Z., 2013. Bioleaching of Pb-Zn-Sn chalcopyrite concentrate in tank bioreactor and microbial community succession analysis. *T Nonferr Metal Soc* 23, 3758-3762.

Wang, S., Zhang, B.G., Diao, M.H., Shi, J.X., Jiang, Y.F., Cheng, Y.T., Liu, H., 2018. Enhancement of synchronous bio-reductions of vanadium (V) and chromium (VI) by mixed anaerobic culture. *Environ Pollut* 242, 249-256.

Warnecke, F., Sommaruga, R., Sekar, R., Hofer, J.S., Pernthaler, J., 2005. Abundances, identity, and growth state of actinobacteria in mountain lakes of different UV transparency. *Appl Environ Microbiol* 71, 5551-5559.

Weinstein, J.N., Myers, T.G., OConnor, P.M., Friend, S.H., Fornace, A.J., Kohn, K.W., Fojo, T., Bates, S.E., Rubinstein, L.V., Anderson, N.L., Buolamwini, J.K., vanOsdol, W.W., Monks, A.P., Scudiero, D.A., Sausville, E.A., Zaharevitz, D.W., Bunow, B., Viswanadhan, V.N., Johnson, G.S., Wittes, R.E., Paull, K.D., 1997. An information-intensive approach to the molecular pharmacology of cancer. *Science* 275, 343-349.

Wen, J.G., Jirka, G.H., Raggio, G., 1994. Linked Sediment/Contaminant Transport Model for Rivers with Application to the Buffalo River, New-York. *J Great Lakes Res* 20, 671-682.

Williams, R.J., Howe, A., Hofmockel, K.S., 2014. Demonstrating microbial co-occurrence pattern analyses within and between ecosystems. *Front Microbiol* 5, 358.

Xi, X.F., Wang, L., Hu, J.J., Tang, Y.S., Hu, Y., Fu, X.H., Sun, Y., Tsang, Y.F., Zhang, Y.N., Chen, J.H., 2014. Salinity influence on soil microbial respiration rate of wetland in the Yangtze River estuary through changing microbial community. *J Environ Sci-China* 26, 2562-2570.

Xu, X.J., Chen, C., Guan, X., Yuan, Y., Wang, A.J., Lee, D.J., Zhang, Z.F., Zhang, J., Zhong, Y.J., Ren, N.Q., 2017. Performance and microbial community analysis of a microaerophilic sulfate and nitrate co-reduction system. *Chemical Engineering Journal* 330, 63-70.

Yan, Z., Hao, Z., Wu, H., Jiang, H., Yang, M., Wang, C., 2018. Co-occurrence patterns of the microbial community in polycyclic aromatic hydrocarbon-contaminated riverine sediments. *J Hazard Mater* 367, 99-108.

Yao, J., Sánchez-Pérez, J.M., Sauvage, S., Teissier, S., Attard, E., Lauga, B., Duran, R., Julien, F., Bernard-Jannin, L., Ramburn, H., Gerino, M., 2017. Biodiversity and ecosystem purification service in an alluvial wetland. *Ecological Engineering* 103, 359-371.

Yergeau, E., Lawrence, J.R., Sanschagrin, S., Waiser, M.J., Korber, D.R., Greer, C.W., 2012. Next-generation sequencing of microbial communities in the Athabasca River and its tributaries in relation to oil sands mining activities. *Appl Environ Microbiol* 78, 7626-7637.

Yin, H., Niu, J., Ren, Y., Cong, J., Zhang, X., Fan, F., Xiao, Y., Zhang, X., Deng, J., Xie, M., He, Z., Zhou, J., Liang, Y., Liu, X., 2015. An integrated insight into the response of sedimentary microbial communities to heavy metal contamination. *Scientific Reports* 5, 14266.

Zaaboub, N., Martins, M.V., Dhib, A., Bejaoui, B., Galgani, F., El Bour, M., Aleya, L., 2015. Accumulation of trace metals in sediments in a Mediterranean Lagoon: Usefulness of metal sediment fractionation and elutriate toxicity assessment. *Environ Pollut* 207, 226-237.

Zapata-Peñasco, I., Salazar-Coria, L., Saucedo-García, M., Villa-Tanaka, L., Hernández-Rodríguez, C., 2013. Bisulfite reductase and nitrogenase genes retrieved from biocorrosive



955 bacteria in saline produced waters of offshore oil recovery facilities. *Int Biodeterior*  
956 *Biodegrad* 81, 17-27.  
957 Zeng, Y., Zou, Y., Chen, B., Grebmeier, J.M., Li, H., Yu, Y., Zheng, T., 2011. Phylogenetic  
958 diversity of sediment bacteria in the northern Bering Sea. *Polar Biology* 34, 907-919.  
959 Zhang, B., Xu, X., Zhu, L., 2017. Structure and function of the microbial consortia of  
960 activated sludge in typical municipal wastewater treatment plants in winter. *Sci Rep* 7, 17930.  
961 Zhang, B., Xu, X., Zhu, L., 2018. Activated sludge bacterial communities of typical  
962 wastewater treatment plants: distinct genera identification and metabolic potential differential  
963 analysis. *AMB Express* 8, 184.

964

965

## Figure legends

**Figure 1. Map of the Ichkeul Lake / Bizerte Lagoon hydrological system.** The sampling stations (ST, red) are indicated by numbers.

**Figure 2. Comparison of bacterial communities from stations of the Ichkeul Lake / Bizerte Lagoon hydrological system.** The canonical correspondence analysis shows the correlations between bacterial communities structures (T-RFLP patterns) and environmental chemical data in a) the Bizerte Lagoon-Ichkeul Lake complex, b) in Ichkeul Lake sediments, and c) in Bizerte Lagoon sediments. The analyses were performed in September 2011 (wet season) and April 2012 (dry season). Lake September, pink circles; Lake April, blue circles; Lagoon September, red diamonds; Lagoon April brown diamonds. The analyses are based on biological triplicates for each station.  $\Sigma$  Pesticides, total pesticides;  $\Sigma$  PAHs, total PAHs;  $\Sigma$  PCBs, total PCB;  $\Sigma$  Sn, total Sn organic compounds; T°, temperature; m-ERM-Q, mean effect range medium quotient; P, phosphate; DO, dissolved oxygen. The names of the stations further characterized in this study are indicated ST5 and ST8 for the Ichkeul Lake, and ST12 and ST17 for the Bizerte Lagoon.

**Figure 3. Composition of bacterial communities from stations of the Ichkeul Lake (ST5 and ST8) and Bizerte Lagoon (ST12 and ST17) hydrological system.** The analysis was performed at the phylum level applying a threshold similarity of 97 % for OTU identification. The average of three replicates was used for the analysis. Other, correspond to phyla with relative abundance < 5%.

**Figure 4. Comparison of bacterial communities from the different stations of the Ichkeul Lake / Bizerte Lagoon hydrological system.** a) Venn diagram showing the number of shared OTUs between stations as well as the specific OTUs for each station. b) Linear discriminant analysis effect size (LEfSe) of the habitat (lake / lagoon). c) LEfSe analysis of the contamination. LEfSe analysis shows the significant differential abundance distribution of bacterial genera between the samples of environmental groups (lake / lagoon; high

contamination / low contamination). The analysis was performed applying a threshold similarity of 97 % for OTU identification. The contamination level (high / low) refers to mean effect range medium quotients (m-ERM-Q) presented in Table 1.

**Figure 5. Heatmap comparing the bacterial communities from the different stations of the Ichkeul Lake / Bizerte Lagoon hydrological system.** The heatmap with clustering analysis based on Pearson correlation shows the repartition of the 50 most abundant genera. The analysis was performed applying a threshold similarity of 97 % for OTU identification. The triplicates for each station are shown in the heatmap.

**Figure 6. Clustered correlation (ClusCor) between bacterial OTUs and pollutants parameters.** The analysis is based on Pearson correlation, red indicating high positive correlation and blue high negative correlation ( $p$  value  $<0.01$ ). Clustering was performed applying Pearson correlations and Ward algorithm as cluster method. Abbreviations of pollutants are those indicated in Table 1. The analysis was performed at the genus level applying a threshold similarity of 97 % for OTU identification. The specific OTUs revealed by LEfSe analysis are coloured according to their specificity: lake (khaki), lagoon (red), high contamination (orange), and low contamination (blue).

**Table 1: Chemical analyses of contaminants in the Ichkeul Lake / Bizerte Lagoon hydrological system and sediment toxicity potential.** The data corresponds to the sum of the different types of pollutants obtained in April 2012 described in details by Ben Salem et al. (2017). Average  $\pm$  SD are presented (n=3).

Pollutants (ng g <sup>-1</sup> dry wt)	Stations			
	ST5 Lake	ST8	ST12 Lagoon	ST17
<b>Organic Sn<sup>a</sup></b>	143 $\pm$ 0.3	83 $\pm$ 0.3	154 $\pm$ 0.1	102 $\pm$ 0.5
<b>PCBs<sup>b</sup></b>	4.66 $\pm$ 0.7 (I) <sup>h</sup>	5.58 $\pm$ 1 (II) <sup>h</sup>	4.54 $\pm$ 0.1 (I) <sup>h</sup>	2.48 $\pm$ 0.8 (I) <sup>h</sup>
<b>Heptachlor</b>	10.7 $\pm$ 0.7	14 $\pm$ 1.6	11 $\pm$ 4	12 $\pm$ 4
<b>Lindane<sup>d</sup></b>	268 $\pm$ 6	283 $\pm$ 13	101 $\pm$ 5	95 $\pm$ 14
<b>Other OCPs<sup>e</sup></b>	586 $\pm$ 0.03 (IV) <sup>h</sup>	1717 $\pm$ 65 (IV) <sup>h</sup>	53 $\pm$ 3.4 (III) <sup>h</sup>	86 $\pm$ 7.4 (III) <sup>h</sup>
<b>PAHs<sup>f</sup></b>	1362 $\pm$ 560 (II) <sup>h</sup>	444 $\pm$ 24 (II) <sup>h</sup>	17988 $\pm$ 1183 (IV) <sup>h</sup>	137 $\pm$ 26 (I) <sup>h</sup>
<b>m-ERM-Q<sup>g</sup></b>	8.73 $\pm$ 0.03	13.30 $\pm$ 0.4	5.30 $\pm$ 0.06	3.90 $\pm$ 0.1

<sup>a</sup>sum of organic forms of tin including methyl, butyl, and phenyl derivatives.

<sup>b</sup>Sum of PCB congeners including PCB18, PCB28, PCB31, PCB44, PCB52, PCB101, PCB138, PCB118, PCB153, PCB149, PCB180, PCB194.

<sup>c</sup>Sum of heptachlor and its epoxide derivative.

<sup>d</sup>Sum of lindane and its metabolites.

<sup>e</sup>Sum of endosulfan, dieldrin, aldrin, chlordane, and their derivatives.

<sup>f</sup>Sum of PAHs including naphthalene, acenaphthylene, acenaphthene, fluorene, phenanthrene, anthracene, fluoranthene, pyrene, benzo[a]anthracene, chrysene, benzo[b]fluoranthene, benzo[k]fluoranthene, benzo[a]pyrene, indeno[1,2,3-cd]pyrene, dibenz[a,h]anthracene, and benzoghiperylene.

<sup>g</sup>m-ERM-Q, mean effect range medium quotient calculated according to (Long et al., 1998).

<sup>h</sup>Quality classification according to Bakke et al. (2010).

1031 **Table 2: Alpha diversity indices (mean  $\pm$  SD, n=3) of microbial community from the**  
1032 **Ichkeul Lake / Bizerte Lagoon stations**

Indices	Station			
	ST5 Lake	ST8 Lake	ST12 Lagoon	ST17 Lagoon
<b>Reads<sup>a</sup></b>	22368	19247	22101	25662
<b>Trimmed sequences<sup>b</sup></b>	16378	14217	15185	18502
<b>Species richness (<i>R</i>)</b>	347 $\pm$ 32	359 $\pm$ 28	341 $\pm$ 27	356 $\pm$ 40
<b>Coverage</b>	0.93 $\pm$ 0.01	0.93 $\pm$ 0.01	0.94 $\pm$ 0.01	0.94 $\pm$ 0.01
<b>Simpson (<i>I-D</i>)</b>	0.97 $\pm$ 0.01	0.98 $\pm$ 0.01	0.96 $\pm$ 0.01	0.98 $\pm$ 0.01
<b>Shannon (<i>H</i>)</b>	4.51 $\pm$ 0.1	4.58 $\pm$ 0.1	4.37 $\pm$ 0.1	4.51 $\pm$ 0.05
<b>Evenness (<i>E</i>)</b>	0.26 $\pm$ 0.04	0.27 $\pm$ 0.01	0.23 $\pm$ 0.03	0.26 $\pm$ 0.02
<b>Chao-1</b>	373 $\pm$ 39	386 $\pm$ 27	362 $\pm$ 27	379 $\pm$ 45

<sup>a</sup>SD: ST5,  $\pm$  1966; ST8,  $\pm$  1866; ST12,  $\pm$  3380; ST17,  $\pm$  1688.

<sup>b</sup>SD: ST5,  $\pm$  1481; ST8,  $\pm$  1406; ST12,  $\pm$  2708; ST17,  $\pm$  1611.















