



HAL
open science

New insights into the biomineralization of mercury selenide nanoparticles through stable isotope analysis in giant petrel tissues

Silvia Queipo-Abad, Zoyne Pedrero, Claudia Marchán-Moreno, Khoulood El Hanafi, Sylvain Bérail, Warren T Corns, Yves Cherel, Paco Bustamante, David Amouroux

► To cite this version:

Silvia Queipo-Abad, Zoyne Pedrero, Claudia Marchán-Moreno, Khoulood El Hanafi, Sylvain Bérail, et al.. New insights into the biomineralization of mercury selenide nanoparticles through stable isotope analysis in giant petrel tissues. *Journal of Hazardous Materials*, 2022, 425, pp.127922. 10.1016/j.jhazmat.2021.127922 . hal-03475304

HAL Id: hal-03475304

<https://univ-pau.hal.science/hal-03475304>

Submitted on 10 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **New insights into the biomineralization of**
2 **mercury selenide nanoparticles through stable**
3 **isotope analysis in giant petrel tissues**

4 **Silvia Queipo-Abad^a, Zoyne Pedrero^{a*}, Claudia Marchán-Moreno^a, Khouloud El**
5 **Hanafi^a, Sylvain Bérail^a, Warren T. Corns^b, Yves Cherel^c, Paco Bustamante^{d,e},**
6 **David Amouroux^a**

7 ^a Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM, Institut des
8 Sciences Analytiques et de Physico-chimie pour l'Environnement et les matériaux, Pau,
9 France.

10 ^b PS Analytical, Arthur House, Crayfields Industrial Estate, Main Road, Orpington, Kent
11 BR5 3HP, UK

12 ^c Centre d'Etudes Biologiques de Chizé, UMR 7372 CNRS - La Rochelle Université,
13 79360 Villiers-en-Bois, France

14 ^d Littoral Environnement et Sociétés (LIENSs), UMR 7266 CNRS - La Rochelle
15 Université, 2 rue Olympe de Gouges, 17000 La Rochelle, France

16 ^e Institut Universitaire de France (IUF), 1 Rue Descartes, 75005 Paris, France

17
18 *Corresponding author: Zoyne Pedrero

19 E-mail address: zoyne.pedrerozayas@univ-pau.fr

20 Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM, Institut des
21 Sciences Analytiques et de Physico-chimie pour l'Environnement et les matériaux, Pau,
22 France

25 **ABSTRACT**

26 Tiemannite (HgSe) is considered the end-product of methylmercury (MeHg) demethylation in
27 vertebrates. The biomineralization of HgSe nanoparticles (NPs) is understood to be an efficient
28 MeHg detoxification mechanism; however, the process has not yet been fully elucidated. In order
29 to contribute to the understanding of complex Hg metabolism and HgSe NPs formation, the Hg
30 isotopic signatures of 40 samples of 11 giant petrels were measured. This seabird species is one
31 of the largest avian scavengers in the Southern Ocean, highly exposed to MeHg through their
32 diet, reaching Hg concentrations in the liver up to more than 900 $\mu\text{g g}^{-1}$. This work constitutes the
33 first species-specific isotopic measurement ($\delta^{202}\text{Hg}$, $\Delta^{199}\text{Hg}$) of HgSe NPs in seabirds and the
34 largest characterization of this compound in biota. Similar $\delta^{202}\text{Hg}$ values specifically associated
35 to HgSe ($\delta^{202}_{\text{HgSe}}$) and tissues ($\delta^{202}_{\text{bulk}}$) dominated by inorganic Hg species were found, suggesting
36 that no isotopic fractionation is induced during the biomineralization step from the precursor
37 (demethylated) species. In contrast, the largest variations between $\delta^{202}\text{Hg}_{\text{bulk}}$ and $\delta^{202}_{\text{HgSe}}$ were
38 observed in muscle and brain tissues. This could be attributed to the higher fraction of Hg present
39 as MeHg in these tissues. Hg-biomolecules screening highlights the importance of the isotopic
40 characterization of these (unknown) complexes.

41 **KEYWORDS**

42 Mercury, seabirds, isotopic fractionation, HgSe nanoparticles, MeHg demethylation

43

44

1 INTRODUCTION

45 Mercury (Hg) is a globally distributed pollutant present in polar and sub-polar areas through long-
46 range atmospheric transport (Fitzgerald et al., 1998). Once in the marine environment, inorganic
47 mercury (iHg) can be methylated by microbial activity to form methylmercury (MeHg) (Fitzgerald
48 et al., 2007; Kaschak et al., 2013; Mason, 2013). MeHg bioaccumulates in the tissues of aquatic
49 organisms, and biomagnifies through the food chain resulting in predators at the top of the trophic
50 chain to accumulate large amounts of Hg in their tissues (Mason et al., 1995).

51 Seabirds in the Southern Hemisphere showed specific patterns of accumulation, distribution and
52 biotransformation of Hg (Albert et al., 2019; Mills et al., 2020; Renedo et al., 2021). Giant petrels
53 are among the top scavengers of these latitudes, feeding on high trophic level species, which
54 leads to the very high concentrations of Hg in their tissues (Renedo et al., 2021). Their great
55 longevity, of up to 50 years old (Foote et al., 2011), coupled with their high trophic level makes
56 them a species of great ecotoxicological interest, as they can bioaccumulate large quantities of
57 toxic elements, especially Hg, which is ingested through the diet as MeHg. Hepatic demethylation
58 is considered a key detoxification pathway of MeHg in seabirds (Kim et al., 1996; Thompson and
59 Furness, 1989) together with its excretion in feathers (Albert et al., 2019). Demethylation involves
60 selenium (Se) (Khan and Wang, 2010), an essential element which presents a high affinity to
61 inorganic mercury (iHg), resulting ultimately in the formation of insoluble tiemannite nanoparticles
62 (HgSe NPs). Although the liver is the key organ for Hg bioaccumulation, HgSe NPs have been
63 found in other organs and tissues of fish, marine mammals and seabirds, including kidneys,
64 muscle and brain (Gajdosechova et al., 2016; Manceau et al., 2021c, 2021a).

65 Hg isotopic analysis in biological samples is a powerful tool to trace Hg trophic sources and
66 metabolic pathways in wildlife (Bolea-Fernandez et al., 2019; Feng et al., 2015; Li et al., 2020;
67 Masbou et al., 2018; Perrot et al., 2016) and in humans (Du et al., 2018; Sherman et al., 2015,
68 2013). Most of these studies have been carried out on mammalian or fish tissues (Bolea-
69 Fernandez et al., 2019; Feng et al., 2015; Le Croizier et al., 2020; Masbou et al., 2018) and only
70 a few of these studies have focused on seabirds (Manceau et al., 2021b; Poulin et al., 2021;
71 Renedo et al., 2021, 2020, 2018a, 2017). Hg stable isotopes can undergo both mass-dependent

72 fractionation (MDF) and mass-independent fractionation (MIF). Odd Hg isotope MIF is related to
73 the processes of photoreduction and photodemethylation of Hg mainly in the photic zone, thus
74 allowing traceable marine food source to prey-predator relationships (Kwon et al., 2013; Le
75 Croizier et al., 2020; Renedo et al., 2018c; Sherman et al., 2013). In contrast, the metabolic
76 distribution of Hg in the body can be traced through MDF (Bolea-Fernandez et al., 2019; Feng et
77 al., 2015). This powerful tool has been successfully exploited to track Hg distribution in the tissues
78 of three different seabirds species, revealing that hepatic MeHg demethylation is followed by
79 internal tissue redistribution of the residual MeHg enriched in heavier Hg isotopes (Renedo et al.,
80 2021).

81 The main aim of this study was to provide new insights into the biomineralization of MeHg into
82 HgSe NPs by the specific isotopic characterization of this inert Hg compound. For this purpose,
83 a large set of giant petrel samples (for a total of 40 tissues) corresponding to 11 individuals of
84 different ages were studied. Complementary analytical techniques have been combined, such as
85 Hg speciation at the chemical and molecular level, as well as Hg and Se measurements between
86 solid and soluble fractions of tissues. The current work represents the first study on Hg isotopic
87 characterization specifically associated to NPs in (sea)birds. Hg isotopic analyses in blood, as
88 well as the comparison between Hg isotopic composition in HgSe NPs and bulk tissues, shedding
89 light about different distribution routes between tissues of seabirds.

90 **2 MATERIAL AND METHODS**

91 **2.1. Ecological characteristics of giant petrels**

92 The avian model corresponds to the two sibling species of giant petrels, the northern
93 (*Macronectes halli*) and southern (*Macronectes giganteus*) giant petrels, which are the dominant
94 scavengers of the Southern Ocean (Rheindt and Austin, 2005; Salomon and Voisin, 2010). Giant
95 petrels are large birds that can reach over two meters of wingspan (Carlos and Voisin, 2008).
96 During the reproduction period, adult males feed mostly on land on carrion, seabirds and
97 pinnipeds, while females mainly forage at sea, preying on fish, squid and krill (Thiers et al., 2014).

98

99 **2.2. Sampling set and individual characteristics**

100 Eleven dead individuals of giant petrels were opportunistically collected. Four of them were *M.*
101 *halli* (all males) and were collected in the Kerguelen archipelago (Southern Indian Ocean) in 2014.
102 The other four birds were *M. giganteus*. One specimen was collected in the Kerguelen
103 Archipelago in 2014 (male) and three were collected in Adélie Land (Antarctic continent) in 2019
104 (1 female and 2 males). The age of these individuals could not be precisely determined, but one
105 of the seabirds collected in Adélie Land was a fully-fledged chick (*viz.* less than 1 year old). In
106 addition to this, samples of three other adults *M. giganteus* (see details in Renedo et al., 2021)
107 from the same region were also considered in the present study. The samples from Adélie Land
108 were in good shape for adults who died after a collision against an electric pole, but the chick (P-
109 P1) was very emaciated suggesting it dies from starvation. For all the others, the cause of death
110 is unknown, in general they were in good condition. All of them were collected freshly after their
111 death. Internal tissues were sampled, weighed, and stored individually in plastic bags. Liver,
112 kidneys and pectoral muscle were collected from all the eight birds, while brain tissue was
113 sampled for the three *M. giganteus* collected in 2019. Blood samples were obtained by collecting
114 clotted blood from heart atria. After dissection, all the samples were stored at -20°C. All details
115 relating to the complete sampling set are provided in Table S1. All the tissues used in this work
116 were homogenized with an ultra-turrax® and a portion was freeze-dried for analytical purposes.

117

118 **2.3. Determination of total Hg, Se and Hg species**

119 **2.3.2. Determination of Hg and Se concentrations**

120 Freeze-dried and homogenized tissue samples (0.05 - 0.10 g) were digested in 3 mL of Trace
121 Metal Grade HNO₃ (Fisher scientific, Illkirch, France) in an Ultrawave microwave digestion system
122 (Milestone, Sorisole, Italy) at 220°C for 10 min. Fresh-frozen blood samples were digested directly
123 without freeze-drying. The measurement of total Hg (THg) in the digests was performed by cold
124 vapor - atomic fluorescence spectrometry (CV-AFS PS Analytical 10.025, Kent, UK) by reduction
125 with 3% (m/v) SnCl₂ (Scharlab, Barcelona, Spain) in 10% (v/v) HCl (J.T.Baker, Fisher scientific,
126 Illkirch, France). The samples were then diluted in HCl 5% (v/v) for analyses and the quantification
127 was performed by standard calibration. For Se analyses, the resulting digests were appropriately
128 diluted with Milli-Q water and analyzed by ICP-MS. Method protocols were validated using
129 certified reference materials (BCR-464, NIST-1947 and DOLT-5). The information relative to the
130 quality assurance of the Hg analyses of Hg and Se is found in Table S2.

131 **2.3.3. Determination of iHg and MeHg species**

132 Between 0.05-0.10 g of freeze-died samples was added to glass vessels together with 3 mL of
133 25% (m/v) tetramethylammonium hydroxide (TMAH) and a magnetic stir bar. For blood, Hg
134 species were directly extracted from the freshly frozen samples. Vials were closed with a Teflon
135 cap and placed in the microwave. The focused microwave assisted extraction of the samples was
136 performed using a Discover SP-Microwave (CEM Corporation, Matthews, NC). The extraction
137 was carried out using a 1-min ramp up to 75°C followed by 3 min at 75°C with constant stirring.
138 Inorganic mercury (iHg) was measured by CV-AFS after dilution of the extract in HCl 5% (v/v)
139 (Davis et al., 2007; Davis and Long, 2011; Renedo et al., 2021). For the determination of total
140 Hg, 50 µL of a 0.1N (0.05 mol/L cBr₂) potassium bromide/bromate solution (Merck, Darmstadt,
141 Germany) was added to 10 mL of the diluted digest in 5% (v/v) HCl. The CV-AFS system was
142 coupled to an oxidation unit (PS Analytical 10.820, Kent, UK) equipped with a UV lamp so that all
143 samples with potassium bromide/bromate were subjected to on-line UV irradiation to facilitate
144 immediate oxidation of MeHg and subsequent measurement of THg. The concentration of MeHg
145 was determined by the subtraction of iHg from the THg concentration (Aranda et al., 2009;
146 Kaercher et al., 2005). This procedure was validated using BCR-464 and TORT-2 as shown in
147 Table S2.

148 **2.4. Determination of Hg biomolecular species by SEC-ICP-MS**

149 The aqueous soluble protein fraction was extracted from fresh samples (approximately 0.1g) by
150 ultra-probe sonication (30 s at 100W power) in 3 mL of 100 mM ammonium acetate (pH 7.4)
151 followed by centrifugation, as described elsewhere (Pedrero et al., 2011). The obtained cytosolic
152 biomolecules were separated by size exclusion chromatography (SEC) using a Superdex 200
153 300/10 column (10mm ID×300mm length ×13 µm particle size) (GE Healthcare, Uppsala,
154 Sweden). A HPLC Agilent 1100 (Agilent, Wilmington, DE) equipped with a binary HPLC pump
155 and an autosampler was coupled to an Agilent ICP-MS 7500ce (Yokogawa Analytical Systems,
156 Tokyo, Japan). The supernatant was analysed by injection of 100 µL fractions. Isocratic elution
157 was performed at 0.7 mL min⁻¹ with a mobile phase of 100 mM ammonium acetate at pH of 7.4
158 (Pedrero et al., 2011; Pedrero Zayas et al., 2014).

159 **2.5. Extraction of HgSe nanoparticles**

160 Isolation of HgSe NPs was carried out by using an adaptation of the method described elsewhere
161 (Bolea-Fernandez et al., 2019; Gajdosechova et al., 2016). In brief, after defatting, a soft acid
162 treatment with formic acid was applied to the samples. The resulting extract was centrifuged by
163 using 50 kDa cut-off filters (Amicon Ultra). The filter was abundantly washed with Milli-Q water
164 until total removal of soluble Hg and Se was achieved (Gajdosechova et al., 2016). Nanoparticles
165 were then recovered by centrifugation for 3 min at 1000 x g. The collected NPs were then digested
166 in 3 mL HNO₃, using the procedure described in section 2.3. This selective extraction protocol is
167 not considered to be quantitative, due to losses in the cut-off filtration steps. The procedure
168 however has been successfully used for Hg isotopic characterization previously since no isotopic
169 fractionation of this solid species is induced during the sample treatment (Bolea-Fernandez et al.,
170 2019).

171 **2.6. Hg stable isotope analysis**

172 Hg isotopic ratios were measured using a multicollector-(MC)-ICP-MS (Nu Instruments,
173 Wrexham, UK) coupled with continuous flow cold vapor generation (CVG) on the digested
174 samples (see above). The isotopes ¹⁹⁸Hg, ¹⁹⁹Hg, ²⁰⁰Hg, ²⁰¹Hg, ²⁰²Hg, ²⁰³Tl, ²⁰⁴Hg, and ²⁰⁵Tl were
175 simultaneously measured in the Faraday cups L2, L1, Ax, H1, H2, H3, H4 and H5, respectively.
176 In order to correct the instrumental mass-bias, NIST SRM 997 thallium standard solution in 2%
177 (v/v) of HNO₃ (Optima grade, Fisher scientific, Illkirch, France) was nebulized continuously
178 through a desolvation unit (DSN) and the calculation was performed using the exponential law
179 model (Bergquist and Blum, 2007; Blum and Bergquist, 2007; Perrot et al., 2010). The sample
180 standard bracketing (SSB) approach was employed for the calculation of isotope ratios relative
181 to the NIST 3133-iHg standard solution. Mass dependent fractionation (MDF) is reported as
182 recommended by Bergquist and Blum (Bergquist and Blum, 2007) relative to the NIST 3133 Hg
183 solution using Eq. 1:

184
$$\delta^{xxx}Hg = \left(\frac{^{xxx/198}Hg_{sample}}{^{xxx/198}Hg_{NIST\ 3133}} - 1 \right) * 1000 \text{ ‰} \quad \text{Eq. 1}$$

185 where xxx is the studied isotope. Mass-independent fractionation (MIF) of Hg is reported using
186 Eq. 2 as the difference between the theoretical value predicted by MDF and the measured values
187 as $\Delta^{199}\text{Hg}$ and $\Delta^{201}\text{Hg}$ in ‰, according to the protocol suggested elsewhere (Blum and Bergquist,
188 2007):

$$189 \quad \Delta_{\nu}^{xxx}\text{Hg} = \delta^{xxx}\text{Hg} - (\delta^{202}\text{Hg} \times \beta_{xxx}) \quad \text{Eq. 2}$$

190 where β_{xxx} is the kinetic mass-dependence scale factor that depends on the isotopes (0.2520
191 for ^{199}Hg , 0.5024 for ^{200}Hg , 0.7520 for ^{201}Hg and 1.493 for ^{204}Hg) (Blum and Bergquist, 2007). All
192 the samples and standards were measured at a concentration of 1 ng g^{-1} of Hg in a 10% (v/v)
193 $\text{HNO}_3/2\%$ (v/v) HCl solution, to avoid any bias caused by differences in concentration. The
194 external reproducibility of the measurements is expressed as $\pm 2\text{SD}$ for 35 measurements of the
195 NIST RM 8610 (former UM Almadén) as 0.09 and 0.05 ‰ for $\delta^{202}\text{Hg}$ and $\Delta^{199}\text{Hg}$, respectively. All
196 the values for the isotopic composition of the NIST RM 8610 and the certified reference materials
197 BCR- 464, NIST 1947 and TORT-2 analysed in this work are presented in Table S3.

198

199 **3 RESULTS AND DISCUSSION**

200 **3.1. Hg concentrations in giant petrel tissues**

201 The total mercury (THg) concentrations in the tissues of giant petrels are among the highest ever
202 reported for seabirds, up to $933 \mu\text{g g}^{-1}$ dry weight (d.w. here and in all reported concentration
203 values) (Table S4). However, THg was highly variable among tissues for each individual and
204 among individuals considering a single tissue. Nonetheless, the liver always showed the highest
205 concentrations among tissues with Hg hepatic concentrations ranging from 99 to $933 \mu\text{g g}^{-1}$ (mean
206 value of 419 ± 317 , $n=9$) for adult giant petrels. These values are in agreement with previously
207 reported values for giant petrels from New Zealand (Stewart et al., 1999). As liver is the main
208 storage organ for Hg in vertebrates, hepatic concentrations provide an indirect indication of the
209 age of animals (Bolea-Fernandez et al., 2019; Gajdosechova et al., 2016). Thus, our sampling
210 covers a large range of ages allowing us to consider if this factor influences the detoxification

211 process of MeHg. The only chick of the study (P-P1) presents the lowest Hg levels in all the tissue
212 types. The Hg level in liver of the chick individual ($1.1 \mu\text{g g}^{-1}$) is consistent by the process of
213 bioaccumulation with age. In kidneys, THg concentrations were higher for most of the adult
214 samples ranging from $7.8\text{-}481 \mu\text{g g}^{-1}$ (mean value of 85 ± 140 , $n=10$) than the already previously
215 reported concentrations for petrels and other large seabirds such as albatrosses (Bocher et al.,
216 2003; Cipro et al., 2014; Kim et al., 1996; Stewart et al., 1999). In muscles, Hg concentrations
217 vary from 1.0 to $52 \mu\text{g g}^{-1}$ (mean value of 18 ± 20 , $n=9$) in adults ($0.10 \mu\text{g g}^{-1}$ for the chick), and
218 are among the highest reported for seabirds (Renedo et al., 2021). In the case of blood, the two
219 adults from the Antarctic continent (P-10, P-11) exhibit concentrations around $0.5 \mu\text{g g}^{-1}$ (wet
220 weight), which was fivefold higher than the concentration found in the chick (P-P1) from the same
221 location. A linear correlation (Pearson's correlation coefficient $\rho= 0.8971$) was found between
222 THg concentrations in liver and muscle (Figure S1), which could indicate that Hg bioaccumulation
223 occurs in both tissues along time, Hg being mainly retained in the liver but its excess being directly
224 transferred towards the muscles as suggested by previous studies (Manceau et al., 2021c;
225 Renedo et al., 2021).

226 Despite the high variability of MeHg content among individuals (Table S4), likely due to their age
227 difference (Gajdosechova et al., 2016), the percentage of organic Hg was higher in brain (34-
228 74%) than in liver and kidneys (2-34%). Regarding muscles, in the adults with lower Hg
229 concentrations (P10 and P11, both from the Antarctic continent), the MeHg fraction reaches up
230 to 71%. In contrast, in the case of individuals showing higher liver THg concentrations, the
231 percentage of MeHg in the muscles was much lower (8-30%), suggesting that the proportion of
232 MeHg decreases with age together with an increase of THg concentrations. Such a trend was
233 already reported for top of the marine food chain marine mammals including seals and toothed
234 whales (Bolea-Fernandez et al., 2019; Dehn et al., 2005; Gajdosechova et al., 2016).

235 **3.2. Hg isotopic composition of tissues: Understanding Hg (re)distribution**

236 The high $\delta^{202}\text{Hg}_{\text{bulk}}$ variability in the different tissues analyzed contrasts with the quite
237 homogenous $\Delta^{199}\text{Hg}_{\text{bulk}}$ values obtained whatever the tissue (Table S5). Thus, MIF ($\Delta^{199}\text{Hg}_{\text{bulk}}$)
238 ranges from 1.13 to 1.73 ‰ among individuals, except in the case of the chick, which present the

239 lowest values of the study. Considering the whole data set, the $\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$ slope is 1.13 ± 0.01
240 ($r = 0.968$), similar to slope values reported for other marine top predators and fish (Figure S2)
241 (Le Croizier et al., 2020; Masbou et al., 2018; Renedo et al., 2018b, 2021). The reference
242 $\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$ slopes for MeHg photodemethylation processes (1.36 ± 0.02) and iHg
243 photoreduction (1.00 ± 0.02) (Bergquist and Blum, 2007) are also presented in Figure S2. The
244 values of $\Delta^{199}\text{Hg}$ for giant petrel tissues agree with those of blood in skua chicks (1.14 ± 0.05 ‰)
245 and subantarctic penguins (1.16 ± 0.05 ‰), which were correlated with latitude (Renedo et al.,
246 2020). Interestingly, MIF values in the single chick collected on the Antarctic continent (P-P1)
247 ranged from 0.88 to 1.1 ‰ for internal tissues, being slightly lower than for the adults. Giant petrel
248 chicks are fed with prey caught near the colony by their parents. Hence, the MIF difference is
249 likely due to its exclusively Antarctic food sources that are characterized by low $\Delta^{199}\text{Hg}$ values
250 (Renedo et al., 2020). In contrast, adults sampled in the Antarctic Continent may reflect an
251 integration of Hg from local prey during the breeding season and Hg prey from lower latitude
252 during the wintering season (Renedo et al., 2021).

253 MDF ($\delta^{202}\text{Hg}_{\text{bulk}}$) values ranged widely from -1.48 up to 3.05 ‰ (Table S5). In all the individuals,
254 the highest $\delta^{202}\text{Hg}$ values correspond to blood samples. The specific differences of $\delta^{202}\text{Hg}$ values
255 between tissues (Figure 1) is attributed to the strong dependence on the ratio of methylated and
256 demethylated Hg (Feng et al., 2015; Perrot et al., 2016; Renedo et al., 2021). MeHg demethylation
257 induces mass dependent fractionation, resulting in an enrichment of the inorganic Hg species in
258 lighter isotopes whilst the residual MeHg is enriched in heavier ones (Perrot et al., 2016;
259 Rodríguez-González et al., 2009). The highest MDF values in blood, also observed in other
260 animal studies, is understood as a consequence of the transport of residual MeHg (enriched in
261 heavier isotopes from incomplete demethylation processes) by the blood stream (Bolea-
262 Fernandez et al., 2019; Li et al., 2020; Ma et al., 2018; Renedo et al., 2021).

263 Regarding brain, it is generally enriched in heavier isotopes ($\delta^{202}\text{Hg}$ 0.03-1.77 ‰) in comparison
264 to liver and kidneys (Table S5), suggesting residual MeHg from the blood streams subsequently
265 accumulate in this organ. In the case of muscles, two different trends are identified (Figure 1,
266 Table S5) according to the hepatic Hg concentration (associated to the age of the animals).

267 Muscles of giant petrel individuals with the highest THg hepatic concentrations (250 - 933 $\mu\text{g g}^{-1}$)
268 are enriched in lighter isotopes ($\delta^{202}\text{Hg}$) in comparison to liver and kidneys, contrasting with
269 muscles of the other group of animals (THg hepatic concentrations lower than 250 $\mu\text{g g}^{-1}$). Even
270 though Hg isotopic characterization in animal's internal tissues is not extensive, the trend
271 observed in the mentioned samples is in agreement with previous studies on marine mammals
272 and seabirds (Bolea-Fernandez et al., 2019; Li et al., 2020; Renedo et al., 2021) and is attributed
273 to the higher proportion of MeHg in muscles than in the liver. As previously discussed, this
274 organomercurial species is expected to be enriched in heavier isotopes compared to the
275 demethylated ones, assuming preferential demethylation of the lighter Hg isotopes (Perrot et al.,
276 2016; Rodríguez-González et al., 2009).

277 Noticeably, the giant petrel individuals with the highest THg hepatic concentrations (oldest ones)
278 exhibit lighter $\delta^{202}\text{Hg}$ ($\delta^{202}\text{Hg}_{\text{bulk}}$) values in muscles than in liver and kidneys (Tables S4 and S5).
279 Similarly, a recent investigation in marine mammals reported a decrease of $\delta^{202}\text{Hg}$ values in the
280 muscles of the oldest group in comparison to the juvenile one (a shift of 0.8 ‰ between both
281 groups) (Bolea-Fernandez et al., 2019). Atypically, muscles of those giant petrels contained
282 mainly iHg (70-93%). This particular enrichment of muscles in demethylated Hg species is the
283 most likely reason why muscles of these old seabirds contain lighter Hg isotopes, highlighting that
284 demethylation occurs effectively in this tissue as previously reported (Manceau et al. 2021c).
285 Furthermore, several pathways and factors could also influence the stable isotope signature. One
286 of them is the MeHg excretion by molting, a well-known excretion strategy in birds (Bearhop et
287 al., 2000) when MeHg is massively remobilized from internal tissues towards the growing feathers
288 (Renedo et al., 2021). Thus, between 70–90% of the Hg load of the birds is excreted into the new
289 plumage during molting (Agusa et al., 2005; Braune, 1987). In addition, the ageing process
290 intensely impacts, among others, muscle metabolism (Burger, 1993; Wone et al., 2018) and could
291 result in MeHg mobilization (Cuvin-Aralar and Furness, 1991). Moreover, specifically in migrating
292 birds, the consumption of energy stores during migratory fasting could contribute to MeHg
293 mobilization and higher MeHg circulating levels (Seewagen et al., 2016). Therefore, the atypical
294 muscle Hg speciation and isotopic composition in the oldest giant petrels could be the result of
295 the different mechanisms mentioned above.

296 3.3. Hg isotopic composition of HgSe nanoparticles (tiemannite)

297 The present work is the first to isolate and measure the Hg isotopic composition of HgSe NPs in
298 seabird tissues. The specific Hg isotopic composition associated to HgSe NPs ($\delta^{202}\text{Hg}_{\text{HgSe}}$) was
299 compared with the bulk ($\delta^{202}\text{Hg}_{\text{bulk}}$) value (Figure 2, Table S5). In general, in individuals exhibiting
300 the highest THg hepatic concentrations ($250\text{-}933\ \mu\text{g g}^{-1}$), a match for both MDF and MIF, was
301 observed in liver and kidneys between THg ($\delta^{202}\text{Hg}_{\text{bulk}}$) and HgSe NPs ($\delta^{202}\text{Hg}_{\text{HgSe}}$) (typically
302 represented in Figure 2.A for seabird P-1). The specific isotopic signature in isolated HgSe NPs
303 found in the muscles of this set of individuals was also quite similar to the isotopic values in the
304 whole tissue. In the other set of individuals (THg hepatic concentrations lower than $250\ \mu\text{g g}^{-1}$),
305 the $\delta^{202}\text{Hg}$ difference between THg and HgSe NPs showed dissimilar trends depending on
306 tissues. Contrary to muscles and brain, most of HgSe NPs in kidneys are enriched in heavier
307 isotopes ($\delta^{202}\text{Hg}$ shift up to $0.56\ \text{‰}$) in comparison to the bulk (Figure 2.D, Table S5) while no
308 tendency is identified in the liver of these individuals. The mentioned pattern, specifically observed
309 in kidneys, could be related to a physiological peculiarity in birds, where a renal portal system
310 allows this organ to be bypassed (Meredith and Johnson-Delaney, 2010). Considering that HgSe
311 NPs is the final product of MeHg breakdown, the observed ^{202}Hg -enrichment of this solid species
312 in liver and kidneys is unexpected. The previous report of Hg isotopic characterization in HgSe is
313 limited to long-finned pilot whales (five livers and two muscles) (Bolea-Fernandez et al., 2019). It
314 is therefore interesting to note a total agreement in the obtained trends in livers and muscles of
315 both animals belonging to two classes of vertebrates.

316 In the giant petrel group exhibiting the highest Hg hepatic concentrations, there is a perfect match
317 between the isotopic signature of THg ($\delta^{202}\text{Hg}_{\text{bulk}}$) and those in HgSe NPs ($\delta^{202}\text{Hg}_{\text{HgSe}}$) for the
318 liver, kidneys and muscles. In those samples, iHg species were the dominant ones (Table S4),
319 with HgSe constituting 95 ± 5 , 61 ± 8 and $35 \pm 15\%$ in liver, kidneys, and muscle, respectively
320 according to X-ray absorption near edge structure (XANES) analyses (Manceau et al., 2021c). In
321 addition to HgSe NPs, another iHg complex exhibiting a molar ratio Hg:Se (1:4) has been reported
322 by these authors. This complex was described as mercury selenocysteinatate, $\text{Hg}(\text{SeCys})_4$ which
323 is considered to be precursor of tiemannite. $\text{Hg}(\text{SeCys})_4$ was found to be present in liver, kidneys,

324 and muscles representing 5 ± 2 , 35 ± 10 and $61 \pm 13\%$ of the THg, respectively (Manceau et al.,
325 2021c). In each tissue, the THg isotopic composition results from the contribution of different Hg
326 species as outlined in Eq 3:

$$327 \quad \delta^{202}\text{Hg}_{\text{bulk}} = f_{\text{MeHg-R}} \times \delta^{202}_{\text{MeHg-R}} + f_{\text{iHg}} \times \delta^{202}_{\text{iHg}} \quad \text{Eq. 3}$$

328 where f is the fraction of each species in the bulk tissue and the contribution of the iHg species
329 could be expressed as follows in Eq 4:

$$330 \quad \delta^{202}\text{Hg}_{\text{iHg}} = f_{\text{HgSe}} \times \delta^{202}_{\text{HgSe}} + f_{\text{Hg:Se (1:4)}} \times \delta^{202}_{\text{Hg:Se (1:4)}} \quad \text{Eq. 4}$$

331 In the described set of samples with the highest Hg concentrations in the liver, the $\delta^{202}\text{Hg}_{\text{bulk}}$ and
332 $\delta^{202}\text{HgSe}$ are identical (Table S5) irrespectively of the proportion of both inorganic species (HgSe
333 and $\text{Hg}(\text{SeCys})_4$). Therefore, we propose that the mentioned inorganic species share a similar
334 $\delta^{202}\text{Hg}$ signature. This finding suggests that no isotopic fractionation seems to be induced during
335 the biomineralization (HgSe NPs) step from the precursor species Hg:Se (1:4).

336 On the other hand, the largest shifts between $\delta^{202}\text{Hg}$ in bulk and the one specifically associated
337 to HgSe NPs are found in muscles and brain (Table S5). The observed variations could be
338 attributed to the higher fraction of Hg present as MeHg in these two tissues. The organomercurial
339 compound is expected to be specifically enriched in heavier isotopes, since MeHg demethylation
340 is followed by internal tissue redistribution of the residual MeHg enriched in heavier Hg isotopes
341 (Bolea-Fernandez et al., 2019; Li et al., 2020; Ma et al., 2018; Renedo et al., 2021).

342 The brain was the organ exhibiting the highest MeHg percentage (Table S4), where HgSe and
343 Hg:Se (1:4) have been also reported (Manceau et al., 2021c). The specific isotopic signature of
344 the organic compound ($\delta^{202}_{\text{MeHg}}$) could be estimated by considering; *i.* the proportion of these
345 inorganic species (Table S4 and XANES data) and *ii.* assuming HgSe and Hg:Se (1:4) share a
346 similar MDF pattern ($\delta^{202}\text{HgSe}$ in Table S5). The obtained $\delta^{202}\text{MeHg}$ values vary between 1.98
347 and 2.94 ‰ for the investigated individuals, confirming an enrichment in lighter isotopes of the
348 demethylated iHg with respect to MeHg (Figure 3). The isotopic measurement in each individual
349 species is undoubtedly the best approach to determine the species-specific isotopic signature.
350 The obtained (estimated) values are in good agreement with the shift (~ 3 ‰ $\delta^{202}\text{Hg}$) between

351 MeHg and iHg species measured by Gas Chromatography coupled to MC-ICPMS, the single
352 study so far that has measured both Hg species in animals (Perrot et al., 2016).

353 Chemical speciation and (total) Hg isotopic analyses have been combined in several recent
354 studies for the estimation of Hg species-specific isotopic signature (Feng et al., 2015; Li et al.,
355 2020; Poulin et al., 2021). The mentioned approach has provided new and valuable information
356 on Hg in biota, however it should be emphasized that Hg speciation has been simplistically
357 reduced to inorganic and MeHg species, without considering the binding of Hg with biomolecules
358 and proteins (Pedrero et al., 2016).

359

360 **3.4. New insights on Se-mediated MeHg breakdown**

361 Despite the great interest related to the understanding of MeHg transformation into the bioinert
362 HgSe NPs (Cid-Barrio et al., 2020), the mechanism is still not fully characterized. According to
363 our results, it seems that the limiting step in the detoxification of MeHg leading to the formation of
364 HgSe NPs is the cleavage of the C-Hg bond. The mentioned MeHg demethylation step is
365 associated with a MDF shift varying between 2 and 3 ‰ (Figure 3), in total agreement with
366 species-specific isotopic values reported in the literature (Perrot et al., 2016; Poulin et al., 2021;
367 Rodríguez-González et al., 2009). This route would be mediated by Se-biomolecules
368 (Asaduzzaman and Schreckenbach, 2011; Khan and Wang, 2010), finally leading to HgSe
369 precipitation in the tissues (Palmisano et al., 1995). The hypothesis that this stage is
370 thermodynamically favorable (Banerjee et al., 2015) could explain the absence of measurable
371 isotopic fractionation between the two inorganic species, the (demethylated) Hg intermediate
372 complex and the HgSe NPs. Deeper investigation focused on Se speciation in giant petrel tissues
373 has revealed that selenoneine is the main Se-compound whatever the considered tissue (authors'
374 unpublished data). The potential involvement of selenoneine in Hg detoxification has been evoked
375 in several works (Achouba et al., 2019; Yamashita et al., 2013). One of the most advanced
376 hypothesis is the mediation of selenoneine in the cleavage of Hg–methyl bonds, based on the
377 demonstration of such capability by 1-methyl-1,3-dihydro-2H-benzimidazole-2-selone,
378 H(sebenzimMe) a structural analogue of the selenoneine (Palmer and Parkin, 2015). The

379 resulting MeHg demethylation compound presents a molar ratio Hg:Se of 1:4, (Palmer and Parkin,
380 2015) as the one reported on giant petrel samples (Manceau et al., 2021c).

381 The complexity of Hg metabolization, including MeHg demethylation, is unquestionable. In
382 general, Hg speciation is mainly limited to the discrimination between inorganic and organic
383 species, with no consideration of Hg linked to proteins and other biomolecules (Pedrero et al.,
384 2016). In the set of samples investigated, a lower Hg proportion was associated to the water-
385 soluble fraction (Table S4). The screening by size exclusion chromatography of the water
386 extractable fraction of internal tissues of giant petrels reveals the presence of several Hg
387 compounds (Figure 4). The diversity of (unknown) Hg species detected in the water-soluble
388 fraction strongly suggests the complexity of Hg metabolic pathways and its interactions with
389 biomolecules in living organisms. The structural identification of Hg metabolites is limited so far
390 to few examples as hemoglobin (-MeHg) and metallothioneins (-iHg) in marine mammals (Pedrero
391 et al., 2011; Pedrero Zayas et al., 2014), peptide complexes in plants (Krupp et al., 2009) and
392 ethylmercury adduct formation of human serum albumin and β -lactoglobulin (Trümpler et al.,
393 2009) due to the great analytical challenges they represent. Despite the significant advances on
394 the understanding of Hg in biota from this and other recent studies (Bolea-Fernandez et al., 2019;
395 Gajdosechova et al., 2016; Manceau et al., 2021c; Pedrero et al., 2012; Pedrero Zayas et al.,
396 2014; Poulin et al., 2021), species-specific isotopic characterization of Hg compounds, including
397 Hg-biomolecules is mandatory to go further on the characterization of Hg metabolic pathways.

398 **4 CONCLUSIONS**

399 In this study, for the first time HgSe NPs species-specific (Hg) isotopic characterization has been
400 carried out in seabirds. The isotopic signature of this compound, which is considered to be the
401 end-product of MeHg demethylation, was determined in numerous tissues including liver, muscle,
402 kidney and brain tissues of individual giant petrels. The Hg isotopic composition matching of HgSe
403 NPs (measured) and other inorganic species (estimated) indicates that no isotopic fractionation
404 appears to be induced during the HgSe NPs biomineralization step from the precursor-
405 demethylated species. The comparison between species-specific Hg isotopic composition of
406 MeHg (estimated) and demethylated species, suggests that a shift of 2-3 ‰ $\delta^{202}\text{Hg}$ could be

407 associated to the cleavage of the C-Hg bond. On the other hand, the screening of water-
408 extractable Hg binding proteins from different tissues reveals the presence of several (unknown)
409 Hg species, probably involved on MeHg detoxification. Advances in analytical chemistry allowing
410 isolation and/or characterization by hyphenated techniques (e.g. HPLC-MC-ICPMS) will be
411 crucial to accurately determine the Hg isotopic composition in these species and understand their
412 role on Hg metabolization.

413 **ACKNOWLEDGEMENTS**

414 The authors thank the Agence Nationale de la Recherche (MERSEL Project No. ANR-18-CE34-
415 0004-01), CNRS (MITI-ISOISEAUX) and the Communauté d'Agglomération Pau Béarn Pyrénées
416 (MERISOTOP) for their financial support. This project has also received funding from the
417 European Union's Horizon 2020 research and innovation programme under the Marie
418 Skłodowska-Curie grant agreement No 101007962". S. Queipo and C. Marchan Moreno
419 acknowledge their postdoctoral and PhD fellowships, respectively, funded by the Region Nouvelle
420 Aquitaine (Project BENESEL). Support was provided to Yves Cherel by the Institut Polaire
421 Français Paul Emile Victor (IPEV programme no. 109, C. Barbraud) and the Terres Australes et
422 Antarctiques Francaises, and to Paco Bustamante by the Institut Universitaire de France (IUF).
423 The authors would like to acknowledge PS Analytical for their instrumental and technical support.

424

425

- 427 Achouba, A., Dumas, P., Ouellet, N., Little, M., Lemire, M., Ayotte, P., 2019. Selenoneine is a
428 major selenium species in beluga skin and red blood cells of Inuit from Nunavik.
429 *Chemosphere* 229, 549–558. <https://doi.org/10.1016/j.chemosphere.2019.04.191>
- 430 Agusa, T., Matsumoto, T., Ikemoto, T., Anan, Y., Kubota, R., Yasunaga, G., Kunito, T., Tanabe,
431 S., Ogi, H., Shibata, Y., 2005. Body distribution of trace elements in black-tailed gulls from
432 Rishiri Island, Japan: Age-dependent accumulation and transfer to feathers and eggs.
433 *Environ. Toxicol. Chem.* 24, 2107–2120. <https://doi.org/10.1897/04-617R.1>
- 434 Albert, C., Renedo, M., Bustamante, P., Fort, J., 2019. Using blood and feathers to investigate
435 large-scale Hg contamination in Arctic seabirds: A review. *Environ. Res.* 177, 108588.
436 <https://doi.org/10.1016/j.envres.2019.108588>
- 437 Aranda, P.R., Pacheco, P.H., Olsina, R.A., Martinez, L.D., Gil, R.A., 2009. Total and inorganic
438 mercury determination in biodiesel by emulsion sample introduction and FI-CV-AFS after
439 multivariate optimization. *J. Anal. At. Spectrom.* 24, 1441–1445.
440 <https://doi.org/10.1039/b903113h>
- 441 Asaduzzaman, A.M., Schreckenbach, G., 2011. Degradation mechanism of methyl mercury
442 selenoamino acid complexes: A computational study. *Inorg. Chem.* 50, 2366–2372.
443 <https://doi.org/10.1021/ic1021406>
- 444 Banerjee, M., Karri, R., Rawat, K.S., Muthuvel, K., Pathak, B., Roy, G., 2015. Chemical
445 Detoxification of Organomercurials. *Angew. Chemie* 127, 9455–9459.
446 <https://doi.org/10.1002/ange.201504413>
- 447 Bearhop, S., Ruxton, G.D., Furness, R.W., 2000. Dynamics of Mercury in Blood and Feathers of
448 Great Skuas. *Environ. Toxicol. Chem.* 19, 1638. [https://doi.org/10.1897/1551-
449 5028\(2000\)019<1638:DOMIBA>2.3.CO;2](https://doi.org/10.1897/1551-5028(2000)019<1638:DOMIBA>2.3.CO;2)
- 450 Bergquist, B.A., Blum, J.D., 2007. Mass-dependent and -independent fractionation of Hg isotopes

451 by photoreduction in aquatic systems. *Science* (80-.). 318, 417–420.
452 <https://doi.org/10.1126/science.1148050>

453 Blum, J.D., Bergquist, B.A., 2007. Reporting of variations in the natural isotopic composition of
454 mercury. *Anal. Bioanal. Chem.* 388, 353–9. <https://doi.org/10.1007/s00216-007-1236-9>

455 Bocher, P., Caurant, F., Miramand, P., Cherel, Y., Bustamante, P., 2003. Influence of the diet on
456 the bioaccumulation of heavy metals in zooplankton-eating petrels at Kerguelen
457 archipelago, Southern Indian Ocean. *Polar Biol.* 26, 759–767.
458 <https://doi.org/10.1007/s00300-003-0552-6>

459 Bolea-Fernandez, E., Rua-Ibarz, A., Krupp, E.M., Feldmann, J., Vanhaecke, F., 2019. High-
460 precision isotopic analysis sheds new light on mercury metabolism in long-finned pilot
461 whales (*Globicephala melas*). *Sci. Rep.* 9, 1–10. <https://doi.org/10.1038/s41598-019-43825->
462 [z](https://doi.org/10.1038/s41598-019-43825-z)

463 Braune, B.M., 1987. Comparison of total mercury levels in relation to diet and molt for nine species
464 of marine birds. *Arch. Environ. Contam. Toxicol.* 16, 217–224.
465 <https://doi.org/10.1007/BF01055802>

466 Burger, J., 1993. Metals in avian feathers: Bioindicators of environmental pollution, Review of
467 environmental toxicology.

468 Carlos, C. J & Voisin, J.-F., 2008. Identifying giant petrels , *Macronectes giganteus* and *M. halli*,
469 in the field and in the hand. *Seabird* 21, 1–15.

470 Cid-Barrio, L., Bouzas-Ramos, D., Salinas-Castillo, A., Ogra, Y., Encinar, J.R., Costa-Fernández,
471 J.M., 2020. Quantitative assessment of cellular uptake and differential toxic effects of HgSe
472 nanoparticles in human cells. *J. Anal. At. Spectrom.* 35, 1979–1988.
473 <https://doi.org/10.1039/d0ja00162g>

474 Cipro, C.V.Z., Cherel, Y., Caurant, F., Miramand, P., Méndez-Fernandez, P., Bustamante, P.,
475 2014. Trace elements in tissues of white-chinned petrels (*Procellaria aequinoctialis*) from

476 Kerguelen waters, Southern Indian Ocean. *Polar Biol.* 37, 763–771.
477 <https://doi.org/10.1007/s00300-014-1476-z>

478 Cuvin-Aralar, M.L.A., Furness, R.W., 1991. Mercury and selenium interaction: A review.
479 *Ecotoxicol. Environ. Saf.* 21, 348–364. [https://doi.org/10.1016/0147-6513\(91\)90074-Y](https://doi.org/10.1016/0147-6513(91)90074-Y)

480 Davis, W.C., Christopher, S.J., Pugh, R.S., Donard, O.F.X., Krupp, E. a, Point, D., Horvat, M.,
481 Gibicar, D., Kljakovic-Gaspic, Z., Porter, B.J., Schantz, M.M., 2007. Certification of
482 methylmercury content in two fresh-frozen reference materials: SRM 1947 Lake Michigan
483 fish tissue and SRM 1974b organics in mussel tissue (*Mytilus edulis*). *Anal. Bioanal. Chem.*
484 387, 2335–41. <https://doi.org/10.1007/s00216-006-1106-x>

485 Davis, W.C., Long, S.E., 2011. Measurements of methylmercury, ethylmercury, and inorganic
486 mercury species in a whole blood standard reference material: SRM 955c—Toxic Elements
487 in Caprine Blood. *J. Anal. At. Spectrom.* 26, 431. <https://doi.org/10.1039/c0ja00175a>

488 Dehn, L.A., Sheffield, G.G., Follmann, E.H., Duffy, L.K., Thomas, D.L., Bratton, G.R., Taylor, R.J.,
489 O'Hara, T.M., 2005. Trace elements in tissues of phocid seals harvested in the Alaskan and
490 Canadian Arctic: Influence of age and feeding ecology. *Can. J. Zool.* 83, 726–746.
491 <https://doi.org/10.1139/z05-053>

492 Du, B., Feng, X., Li, P., Yin, R., Yu, B., Sonke, J.E., Guinot, B., Anderson, C.W.N., Maurice, L.,
493 2018. Use of Mercury Isotopes to Quantify Mercury Exposure Sources in Inland Populations,
494 China. *Environ. Sci. Technol.* 52, 5407–5416. <https://doi.org/10.1021/acs.est.7b05638>

495 Feng, C., Pedrero, Z., Gentès, S., Barre, J., Renedo, M., Tessier, E., Berail, S., Maury-Brachet,
496 R., Mesmer-Dudons, N., Baudrimont, M., Legeay, A., Maurice, L., Gonzalez, P., Amouroux,
497 D., 2015. Specific Pathways of Dietary Methylmercury and Inorganic Mercury Determined
498 by Mercury Speciation and Isotopic Composition in Zebrafish (*Danio rerio*). *Environ. Sci.*
499 *Technol.* 49, 12984–12993. <https://doi.org/10.1021/acs.est.5b03587>

500 Fitzgerald, W.F., Engstrom, D.R., Mason, R.P., Nater, E.A., 1998. The case for atmospheric
501 mercury contamination in remote areas. *Environ. Sci. Technol.* 32, 1–7.

502 <https://doi.org/10.1021/es970284w>

503 Fitzgerald, W.F., Lamborg, C.H., Hammerschmidt, C.R., 2007. Marine Biogeochemical Cycling of
504 Mercury. *Chem. Rev.* 107, 641–662. <https://doi.org/10.1021/cr050353m>

505 Foote, C.G., Daunt, F., González-Solís, J., Nasir, L., Phillips, R.A., Monaghan, P., 2011. Individual
506 state and survival prospects: age, sex, and telomere length in a long-lived seabird. *Behav.*
507 *Ecol.* 22, 156–161. <https://doi.org/10.1093/beheco/arq178>

508 Gajdosechova, Z., Lawan, M.M., Urgast, D.S., Raab, A., Scheckel, K.G., Lombi, E., Kopittke,
509 P.M., Loeschner, K., Larsen, E.H., Woods, G., Brownlow, A., Read, F.L., Feldmann, J.,
510 Krupp, E.M., 2016. In vivo formation of natural HgSe nanoparticles in the liver and brain of
511 pilot whales. *Sci. Rep.* 6, 1–11. <https://doi.org/10.1038/srep34361>

512 Kaercher, L.E., Goldschmidt, F., Paniz, J.N.G., De Moraes Flores, É.M., Dressler, V.L., 2005.
513 Determination of inorganic and total mercury by vapor generation atomic absorption
514 spectrometry using different temperatures of the measurement cell. *Spectrochim. Acta - Part*
515 *B At. Spectrosc.* 60, 705–710. <https://doi.org/10.1016/j.sab.2005.03.006>

516 Kaschak, E., Knopf, B., Petersen, J.H., Bings, N.H., König, H., 2013. Biotic methylation of mercury
517 by intestinal and sulfate-reducing bacteria and their potential role in mercury accumulation
518 in the tissue of the soil-living *Eisenia foetida*. *Soil Biol. Biochem.* 69, 1–10.
519 <https://doi.org/10.1016/j.soilbio.2013.11.004>

520 Khan, M.A.K., Wang, F., 2010. Chemical demethylation of methylmercury by selenoamino acids.
521 *Chem. Res. Toxicol.* 23, 1202–1206. <https://doi.org/10.1021/tx100080s>

522 Kim, E.Y., Murakami, T., Saeki, K., Tatsukawa, R., 1996. Environmental Contamination and
523 Toxicology Mercury Levels and its Chemical Form in Tissues and Organs of Seabirds. *Arch.*
524 *Environ. Contam. Toxicol.* 30, 259–266. <https://doi.org/10.1007/BF00215806>

525 Kritee, K., Barkay, T., Blum, J.D., 2009. Mass dependent stable isotope fractionation of mercury
526 during mer mediated microbial degradation of monomethylmercury. *Geochim. Cosmochim.*

527 Acta 73, 1285–1296. <https://doi.org/10.1016/j.gca.2008.11.038>

528 Kritee, K., Blum, J.D., Johnson, M.W., Bergquist, B.A., Barkay, T., 2007. Mercury stable isotope
529 fractionation during reduction of Hg(II) to Hg(0) by Mercury resistant microorganisms.
530 Environ. Sci. Technol. 41, 1889–1895. <https://doi.org/10.1021/es062019t>

531 Krupp, E.M., Mestrot, A., Wielgus, J., Meharg, A.A., Feldmann, J., 2009. The molecular form of
532 mercury in biota: Identification of novel mercury peptide complexes in plants. Chem.
533 Commun. 28, 4257–4259. <https://doi.org/10.1039/b823121d>

534 Kwon, S.Y., Blum, J.D., Chirby, M.A., Chesney, E.J., 2013. Application of mercury isotopes for
535 tracing trophic transfer and internal distribution of mercury in marine fish feeding
536 experiments. Environ. Toxicol. Chem. 32, 2322–2330. <https://doi.org/10.1002/etc.2313>

537 Le Croizier, G., Lorrain, A., Sonke, J.E., Jaquemet, S., Schaal, G., Renedo, M., Besnard, L.,
538 Cherel, Y., Point, D., 2020. Mercury isotopes as tracers of ecology and metabolism in two
539 sympatric shark species. Environ. Pollut. 265, 114931.
540 <https://doi.org/10.1016/j.envpol.2020.114931>

541 Li, M., Juang, C.A., Ewald, J.D., Yin, R., Mikkelsen, B., Krabbenhoft, D.P., Balcom, P.H.,
542 Dassuncao, C., Sunderland, E.M., 2020. Selenium and stable mercury isotopes provide new
543 insights into mercury toxicokinetics in pilot whales. Sci. Total Environ. 710, 136325.
544 <https://doi.org/10.1016/j.scitotenv.2019.136325>

545 Ma, L., Evans, R.D., Wang, W., Georg, R.B., 2018. In vivo fractionation of mercury isotopes in
546 tissues of a mammalian carnivore (*Neovison vison*). Sci. Total Environ. 627, 1228–1233.
547 <https://doi.org/10.1016/j.scitotenv.2018.01.296>

548 Manceau, A., Azemard, S., Hédouin, L., Vassileva, E., Lecchini, D., Fauvelot, C., Swarzenski,
549 P.W., Glatzel, P., Bustamante, P., Metian, M., 2021a. Chemical forms of mercury in blue
550 marlin billfish: Implications for human exposure. Environ. Sci. Technol. Lett. 8, 405–411.
551 <https://doi.org/10.1021/acs.estlett.1c00217>

552 Manceau, A., Brossier, R., Janssen, S.E., Rosera, T.J., Krabbenhoft, D.P., Cherel, Y.,
553 Bustamante, P., Poulin, B.A., 2021b. Mercury Isotope Fractionation by Internal
554 Demethylation and Biomineralization Reactions in Seabirds: Implications for Environmental
555 Mercury Science. *Environ. Sci. Technol.* <https://doi.org/10.1021/acs.est.1c04388>

556 Manceau, A., Gaillot, A.C., Glatzel, P., Cherel, Y., Bustamante, P., 2021c. In Vivo Formation of
557 HgSe Nanoparticles and Hg-Tetraselenolate Complex from Methylmercury in Seabirds -
558 Implications for the Hg-Se Antagonism. *Environ. Sci. Technol.* 55, 1515–1526.
559 <https://doi.org/10.1021/acs.est.0c06269>

560 Masbou, J., Sonke, J.E., Amouroux, D., Guillou, G., Becker, P.R., Point, D., 2018. Hg-Stable
561 Isotope Variations in Marine Top Predators of the Western Arctic Ocean. *ACS Earth Sp.*
562 *Chem.* 2, 479–490. <https://doi.org/10.1021/acsearthspacechem.8b00017>

563 Mason, R.P., 2013. The Methylation of Metals and Metalloids in Aquatic Systems, in: *Methylation-*
564 *From DNA, RNA and Histones to Diseases and Treatment.* Intech, pp. 271–301.

565 Mason, R.P., Reinfelder, J.R., Morel, F.M.M., 1995. Bioaccumulation of mercury and
566 methylmercury. *Water, Air, Soil Pollut.* 80, 915–921. <https://doi.org/10.1007/BF01189744>

567 Meng, M., Sun, R., Liu, H., Yu, B., Yin, Y., Hu, L., Chen, J., Shi, J., Jiang, G., 2020. Mercury
568 isotope variations within the marine food web of Chinese Bohai Sea: Implications for
569 mercury sources and biogeochemical cycling. *J. Hazard. Mater.* 384, 121379.
570 <https://doi.org/10.1016/j.jhazmat.2019.121379>

571 Meredith, A., Johnson-Delaney, C., 2010. *BSAVA Manual of Exotic Pets*, 5th Editio. ed. British
572 Small Animal Veterinary Association, Quedgeley.

573 Mills, W.F., Bustamante, P., McGill, R.A.R., Anderson, O.R.J., Bearhop, S., Cherel, Y., Votier,
574 S.C., Phillips, R.A., 2020. Mercury exposure in an endangered seabird: Long-term changes
575 and relationships with trophic ecology and breeding success: Mercury exposure in
576 albatrosses. *Proc. R. Soc. B Biol. Sci.* 287, 20202683.
577 <https://doi.org/10.1098/rspb.2020.2683>

578 Palmer, J.H., Parkin, G., 2015. Protolytic Cleavage of Hg-C Bonds Induced by 1-Methyl-1,3-
579 dihydro-2 H -benzimidazole-2-selone: Synthesis and Structural Characterization of Mercury
580 Complexes. *J. Am. Chem. Soc.* 137, 4503–4516. <https://doi.org/10.1021/jacs.5b00840>

581 Palmisano, F., Cardellicchio, N., Zambonin, P.G., 1995. Speciation of mercury in dolphin liver: A
582 two-stage mechanism for the demethylation accumulation process and role of selenium.
583 *Mar. Environ. Res.* 40, 109–121. [https://doi.org/10.1016/0141-1136\(94\)00142-C](https://doi.org/10.1016/0141-1136(94)00142-C)

584 Pedrero, Z., Donard, O.F.X., Amouroux, D., 2016. Pushing back the frontiers of mercury
585 speciation using a combination of biomolecular and isotopic signatures : challenge and
586 perspectives. *Anal. Bioanal. Chem.* 408, 2641–8. [https://doi.org/10.1007/s00216-015-9243-](https://doi.org/10.1007/s00216-015-9243-8)
587 8

588 Pedrero, Z., Mounicou, S., Monperrus, M., Amouroux, D., 2011. Investigation of Hg species
589 binding biomolecules in dolphin liver combining GC and LC-ICP-MS with isotopic tracers. *J.*
590 *Anal. At. Spectrom.* 26, 187–194. <https://doi.org/10.1039/c0ja00154f>

591 Pedrero, Z., Ouerdane, L., Mounicou, S., Lobinski, R., Monperrus, M., Amouroux, D., 2012.
592 Identification of mercury and other metals complexes with metallothioneins in dolphin liver
593 by hydrophilic interaction liquid chromatography with the parallel detection by ICP MS and
594 electrospray hybrid linear/orbital trap MS/MS. *Metallomics* 4, 473–479.
595 <https://doi.org/10.1039/c2mt00006g>

596 Pedrero Zayas, Z., Ouerdane, L., Mounicou, S., Lobinski, R., Monperrus, M., Amouroux, D., 2014.
597 Hemoglobin as a major binding protein for methylmercury in white-sided dolphin liver. *Anal.*
598 *Bioanal. Chem.* 406, 1121–1129. <https://doi.org/10.1007/s00216-013-7274-6>

599 Perrot, V., Epov, V.N., Valentina, M.V.P., Grebenshchikova, I., Zouiten, C., Sonke, J.E., Husted,
600 S., Donard, O.F.X., Amouroux, D., 2010. Tracing sources and bioaccumulation of mercury
601 in fish of Lake Baikal - Angara River using Hg isotopic composition. *Environ. Sci. Technol.*
602 44, 8030–8037. <https://doi.org/10.1021/es101898e>

603 Perrot, V., Masbou, J., Pastukhov, M. V, Epov, V.N., Point, D., Bérail, S., Becker, P.R., Sonke,

604 J.E., Amouroux, D., 2016. Natural Hg isotopic composition of different Hg compounds in
605 mammal tissues as a proxy for in vivo breakdown of toxic methylmercury. *Metallomics* 8,
606 170–178. <https://doi.org/10.1039/C5MT00286A>

607 Poulin, B.A., Janssen, S.E., Rosera, T.J., Krabbenhoft, D.P., Eagles-smith, C.A., Ackerman, J.T.,
608 Stewart, A.R., Kim, E., Kim, J., Manceau, A., 2021. Isotope Fractionation from In Vivo
609 Methylmercury Detoxification in Waterbirds. *ACS Earth Sp. Chem.* 5, 990–997.
610 <https://doi.org/10.1021/acsearthspacechem.1c00051>

611 Renedo, M., Amouroux, D., Duval, B., Carravieri, A., Tessier, E., Barre, J., Bérail, S., Pedrero, Z.,
612 Cherel, Y., Bustamante, P., 2018a. Seabird Tissues As Efficient Biomonitoring Tools for Hg
613 Isotopic Investigations: Implications of Using Blood and Feathers from Chicks and Adults.
614 *Environ. Sci. Technol.* 52, 4227–4234. <https://doi.org/10.1021/acs.est.8b00422>

615 Renedo, M., Amouroux, D., Duval, B., Carravieri, A., Tessier, E., Barre, J., Bérail, S., Pedrero, Z.,
616 Cherel, Y., Bustamante, P., 2018b. Seabird Tissues As Efficient Biomonitoring Tools for Hg
617 Isotopic Investigations: Implications of Using Blood and Feathers from Chicks and Adults.
618 *Environ. Sci. Technol.* 52, 4227–4234. <https://doi.org/10.1021/acs.est.8b00422>

619 Renedo, M., Amouroux, D., Pedrero, Z., Bustamante, P., Cherel, Y., 2018c. Identification of
620 sources and bioaccumulation pathways of MeHg in subantarctic penguins: A stable isotopic
621 investigation. *Sci. Rep.* 8, 1–10. <https://doi.org/10.1038/s41598-018-27079-9>

622 Renedo, M., Bustamante, P., Cherel, Y., Pedrero, Z., Tessier, E., Amouroux, D., 2020. A “seabird-
623 eye” on mercury stable isotopes and cycling in the Southern Ocean. *Sci. Total Environ.* 742.
624 <https://doi.org/10.1016/j.scitotenv.2020.140499>

625 Renedo, M., Bustamante, P., Tessier, E., Pedrero, Z., Cherel, Y., Amouroux, D., 2017.
626 Assessment of mercury speciation in feathers using species-specific isotope dilution
627 analysis. *Talanta* 174, 100–110. <https://doi.org/10.1016/j.talanta.2017.05.081>

628 Renedo, M., Pedrero, Z., Amouroux, D., Cherel, Y., Bustamante, P., 2021. Mercury isotopes of
629 key tissues document mercury metabolic processes in seabirds. *Chemosphere* 263,

630 127777. <https://doi.org/10.1016/j.chemosphere.2020.127777>

631 Rheindt, F.E., Austin, J.J., 2005. Major analytical and conceptual shortcomings in a recent
632 taxonomic revision of the Procellariiformes - A reply to Penhallurick and Wink (2004). *Emu*
633 105, 181–186. <https://doi.org/10.1071/MU04039>

634 Rodríguez-González, P., Epov, V.N., Bridou, R., Tessier, E., Guyoneaud, R., Monperrus, M.,
635 Amouroux, D., 2009. Species-specific stable isotope fractionation of mercury during Hg(II)
636 methylation by an anaerobic bacteria (*Desulfobulbus propionicus*) under dark conditions.
637 *Environ. Sci. Technol.* 43, 9183–8. <https://doi.org/10.1021/es902206j>

638 Salomon, M., Voisin, J.F., 2010. Ecogeographical variation in the southern giant petrel
639 (*Macronectes giganteus*). *Can. J. Zool.* 88, 195–203. <https://doi.org/10.1139/Z09-134>

640 Seewagen, C.L., Cristol, D.A., Gerson, A.R., 2016. Mobilization of mercury from lean tissues
641 during simulated migratory fasting in a model songbird. *Sci. Rep.* 6, 6–10.
642 <https://doi.org/10.1038/srep25762>

643 Sherman, L.S., Blum, J.D., Basu, N., Rajaei, M., Evers, D.C., Buck, D.G., Petrlik, J., DiGangi, J.,
644 2015. Assessment of mercury exposure among small-scale gold miners using mercury
645 stable isotopes. *Environ. Res.* 137, 226–234. <https://doi.org/10.1016/j.envres.2014.12.021>

646 Sherman, L.S., Blum, J.D., Franzblau, A., Basu, N., 2013. New insight into biomarkers of human
647 mercury exposure using naturally occurring mercury stable isotopes. *Environ. Sci. Technol.*
648 47, 3403–3409. <https://doi.org/10.1021/es305250z>

649 Stewart, F.M., Phillips, R.A., Bartle, A.J., Craig, J., Shooter, D., 1999. Influence of phylogeny,
650 diet, moult schedule and sex on heavy metal concentrations in New Zealand
651 Procellariiformes. *Mar. Ecol. Prog. Ser.* 178, 295–305. <https://doi.org/10.3354/meps178295>

652 Thiers, L., Delord, K., Barbraud, C., Phillips, R.A., Pinaud, D., Weimerskirch, H., 2014. Foraging
653 zones of the two sibling species of giant petrels in the Indian Ocean throughout the annual
654 cycle: Implication for their conservation. *Mar. Ecol. Prog. Ser.* 499, 233–248.

655 <https://doi.org/10.3354/meps10620>

656 Thompson, D.R., Furness, R.W., 1989. The chemical form of mercury stored in South Atlantic
657 seabirds. *Environ. Pollut.* 60, 305–317. [https://doi.org/10.1016/0269-7491\(89\)90111-5](https://doi.org/10.1016/0269-7491(89)90111-5)

658 Trümpler, S., Lohmann, W., Meermann, B., Buscher, W., Sperling, M., Karst, U., 2009. Interaction
659 of thimerosal with proteins - Ethylmercury adduct formation of human serum albumin and β -
660 lactoglobulin A. *Metallomics* 1, 87–91. <https://doi.org/10.1039/b815978e>

661 Wone, B.W.M., Kinchen, J.M., Kaup, E.R., Wone, B., 2018. A procession of metabolic alterations
662 accompanying muscle senescence in *Manduca sexta*. *Sci. Rep.* 8, 1–13.
663 <https://doi.org/10.1038/s41598-018-19630-5>

664 Yamashita, M., Yamashita, Y., Suzuki, T., Kani, Y., Mizusawa, N., Imamura, S., Takemoto, K.,
665 Hara, T., Hossain, M.A., Yabu, T., Touhata, K., 2013. Selenoneine, a Novel Selenium-
666 Containing Compound, Mediates Detoxification Mechanisms against Methylmercury
667 Accumulation and Toxicity in Zebrafish Embryo. *Mar. Biotechnol.* 15, 559–570.
668 <https://doi.org/10.1007/s10126-013-9508-1>

669

670

Figure captions

671 **Figure 1.** $\delta^{202}\text{Hg}$ plotted against the percentage of MeHg for the 41 tissue samples of giant
672 petrels. Grey markers correspond to isotopic values reported elsewhere (Renedo et al., 2021) in
673 a pilot study for three giant petrels (PGA01, PGA02 and PGA03). White-filled markers correspond
674 to isotopic values of the youngest individual (P-P1).

675

676 **Figure 2.** $\Delta^{199}\text{Hg}$ vs $\delta^{202}\text{Hg}$ values for liver, kidneys and muscle of giant petrels: A) P-1, B) P-11,
677 C) P-P1 and D) P-10. Dotted arrows correspond to the shift between $\delta^{202}\text{Hg}$ values from THg to
678 HgSe NPs in different organs. Uncertainties in the measurement of the isotopic ratios are
679 expressed as 2SD on the secondary standard RM 8610.

680

681 **Figure 3.** Simplified diagram for the estimation of MDF ($\delta^{202}\text{Hg}$) values of MeHg according to the
682 new insights found for the demethylation/biomineralization of MeHg into HgSe NPs in brain of
683 three giant petrels (PGA01, PGA02 and PGA03).

684

685 **Figure 4.** Typical ^{202}Hg chromatogram obtained by SEC-ICP-MS for the water-soluble fraction of:
686 A) liver and B) kidneys of adult giant petrels.