



HAL
open science

Selenium distribution and speciation in waters of pristine alpine lakes from central-western Pyrenees (France–Spain)

Maïté Bueno, Bastien Duval, Emmanuel Tessier, Andrea Romero-Rama, Leire Kortazar, Luís Ángel Fernández, Alberto de Diego, David Amouroux

► **To cite this version:**

Maïté Bueno, Bastien Duval, Emmanuel Tessier, Andrea Romero-Rama, Leire Kortazar, et al.. Selenium distribution and speciation in waters of pristine alpine lakes from central-western Pyrenees (France–Spain). *Environmental Science: Processes & Impacts*, 2022, 24 (9), pp.1430-1442. 10.1039/D1EM00430A . hal-03807710

HAL Id: hal-03807710

<https://univ-pau.hal.science/hal-03807710v1>

Submitted on 21 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Selenium distribution and speciation in waters of pristine alpine lakes from central-western**
2 **Pyrenees (France-Spain)**

3
4 Maité Bueno^{1*}, Bastien Duval^{1,2}, Emmanuel Tessier¹, Andrea Romero-Rama¹, Leire Kortazar², Luís
5 Ángel Fernández^{2,3}, Alberto de Diego^{2,3}, David Amouroux^{1*}

6
7 ¹ Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, Institute of Analytical Sciences and
8 Physical-Chemistry for the Environment and Materials - IPREM, Pau, France

9 ² Department of Analytical Chemistry, Faculty of Science and Technology, University of the Basque
10 Country (UPV/EHU), Sarriena Auzoa z/g, 48940 Leioa, Basque Country, Spain

11 ³ Plentziako Itsas Estazioa (PIE), University of the Basque Country (UPV/EHU), Areatza
12 pasealekua, 48620 Plentzia, Basque Country, Spain

13

14 Corresponding authors: D Amouroux, M. Bueno

15

16 **Abstract**

17 The speciation of both redox reactive and volatile selenium (Se) compounds, barely reported in
18 pristine aquatic environments, has never been investigated in remote alpine lakes, considered as
19 sensitive ecosystems to detect the effect of global change. This work presents an integrated
20 investigation on Se distribution and speciation conducted in 20 high altitude pristine lakes from the
21 central-western Pyrenees. Five seasonal sampling campaigns were carried out after snowmelt
22 (June/July) and in early fall (October) for the period 2017–2019. Concentrations of total dissolved
23 Se (TDSe) ranged from 7 to 78 ng L⁻¹, being selenate ubiquitously observed in most cases (median
24 of 61% of TDSe). Selenite was only occasionally detected up to 4 ng L⁻¹, therefore a fraction of
25 TDSe was presumably in the forms of elemental Se(0) and/or selenides. Depth profiles carried out
26 in different lakes showed the occurrence of such Se(-II, 0) pool in bottom hypoxic to anoxic waters.
27 The production of volatile Se compounds presented a low median total concentration (TVSe) of 33
28 pg L⁻¹ (range 3–120 pg L⁻¹), mainly in the form of dimethylselenide in sub-surface samples (median
29 of 82% of TVSe). Se concentration in lake waters was significantly correlated with sulphate
30 concentration ($\rho=0.93$, $p<0.0001$), demonstrating that it is influenced by erosion and dissolution of
31 Se and S-enriched parent bedrocks. In addition, for Se depleted alpine lake-bedrock systems, long-
32 range transport and wet atmospheric depositions represent a major source of Se for lake waters.

33

34

35 **1. Introduction**

36 High-mountain lakes are pristine ecosystems characterized by extreme environmental conditions.
37 No data on selenium behaviour have been reported in these aquatic environments. The existing data
38 about Se biogeochemistry in lakes are mainly provided by research in low altitude temperate and
39 tropical lakes and have been reviewed elsewhere (Conde & Sanz Alaejos, 1997; Simmons &
40 Wallschläger, 2005; Ponton et al., 2020). In unpolluted waters, total Se concentrations are typically
41 below 200 ng L⁻¹ (Ponton et al., 2020). In more extreme environments, reported total Se
42 concentrations in western Siberian lakes range from 12 to 311 ng L⁻¹ (Pokrovsky et al., 2018). In
43 Canadian subarctic thermokarst ponds, total dissolved Se ranged from 18 to 67 ng L⁻¹ (Lanceleur et
44 al., 2019), similarly to values from 20 to 60 ng L⁻¹ reported in Norwegian (Økelsrud et al., 2016) or
45 Finish (Wang et al., 1994; 1995) lakes. In North Carolina, Se concentrations in lakes impacted by
46 coal combustion effluents were between 0.3 and 2 µg L⁻¹ while Se was not detected (< 0.22 µg L⁻¹)
47 in reference lakes (Brandt et al., 2017). Among its four oxidation states (-II, 0, IV and VI), Se
48 oxyanions, particularly selenate, are expected to be the major species in well-aerated surface waters
49 and oligotrophic environments (Sharma et al., 2014). Aqueous selenium speciation measurements
50 in lake waters indicated the predominance of selenite (Robberecht et al., 1983; Ponton & Hare,
51 2013; Tanzer & Heumann, 1991) or selenate (Lindström, 1980; Nishri et al., 1999; Ponton & Hare,
52 2013). Despite the lack of analytical procedures to identify and quantify elemental selenium and
53 selenide forms, several authors have reported the presence of such Se pool in lake waters. For
54 example in Finish lakes, the sum of selenite and selenate represented less than 20% of total Se
55 (Wang et al., 1994; 1995). In Lake Kinneret up to 40% of total Se was associated with colloidal
56 elemental Se and selenide forms (Nishri et al., 1999). In Canadian lakes, organo-Se proportion could
57 attain 95% of total Se (Ponton & Hare, 2013).

58 Reduced Se include also volatile selenides seldom tested and reported in lake waters. Reported total
59 volatile Se concentrations varied in a wide range (1 pg L⁻¹ to 109·ng L⁻¹), being dimethylselenide

60 (DMSe) the major species ($\geq 49\%$), followed by dimethyl selenide sulphide (DMSeS) (4–41%) and
61 dimethyl diselenide (DMDS₂) ($\leq 7\%$) (Lanceleur et al., 2019 and references therein). In a mountain
62 minerotrophic peatland (elevation 972 m a.s.l.) a DMSe concentration of 10 ng L⁻¹ was reported in
63 surface water samples (Vriens et al., 2014). In Canadian thermokarst ponds, Lanceleur et al. (2019)
64 found DMSe to be the main volatile species with concentrations between <0.17 to 31 pg Se L⁻¹,
65 while in European estuaries, volatile Se ranged from 22–8067 pg L⁻¹, being DMSe the main species
66 found (Amouroux & Donard, 1997; Tessier et al., 2002).

67 Ninety two per cent of Pyrenean lakes are classified as ultraoligotrophic or oligotrophic (Catalan et
68 al., 2006). Bedrock composition associated with slow weathering and limited soil development
69 determine waters with low mineralization (Catalan et al., 2006). In high altitude mountain lakes, Se
70 can originate either from local (geogenic) or remote sources (atmospheric transport). In the
71 Pyrenees, the main groups of rocks are granites, sedimentary limestones, metamorphic slates and
72 schists, detrital sandstones (Catalan et al., 2006). These rocks generally contain low selenium
73 concentrations (0.01 – 0.08 mg kg⁻¹) although shales can show variable concentrations worldwide
74 (0.1 – 675 mg kg⁻¹) (Plant et al., 2014; Fernández-Martínez & Charlet, 2009). Previous studies have
75 estimated that Se wet deposition (7.4–20.0 Gg Se yr⁻¹) was responsible for 80% of its global
76 atmospheric deposition and was thus dominant over its dry deposition (1.1–5.0 Gg Se yr⁻¹) (Blazina
77 et al., 2017; Pan & Wang, 2015). Suess et al. (2019) reported Se concentration values in rain samples
78 from Pic du Midi (central Pyrenees) between 13–184 ng L⁻¹. The authors determined that the
79 dominant source of Se in rain at the Pic du Midi was the North Atlantic Ocean (54%) with a
80 contribution of terrestrial emissions of 40% with seasonal fluctuation.

81 Pyrenean lakes are also characterized by surface frost periods comprised generally between
82 November and May (Gascoin et al., 2015; López-Moreno & Vicente-Serrano, 2007). During these
83 ice-covered periods, high mountain lakes have shown oxygen depletion leading to anoxic conditions
84 in bottom waters (Catalan et al., 2002). These winter conditions, when oxygen is consumed, may

85 promote conditions favourable to microbial (and possibly sequential) reduction of Se oxyanions to
86 elemental Se, process which has been shown to be environmentally significant (Nancharaiah &
87 Lens, 2015) and may lead to selenium settle out of water column. Although situated in different
88 climatic conditions, depletion of total dissolved selenium and selenate concentrations were
89 previously observed in Lake Kinneret bottom waters when O₂ and SO₄²⁻ were consumed (Nishri et
90 al., 1999; Be'eri-Shlevin et al., 2021).

91 In this study, we surveyed during three years the seasonal concentrations of total Se and its species
92 (including non-volatile and volatile dissolved species) in twenty alpine lakes of the central-western
93 Pyrenees. To this end, sub-surface water samples were collected in June/July 2017–2019 (late
94 spring–early summer) and October 2017–2018 (early fall). Additionally, depth-profiles and
95 temporal monitoring were conducted at some lakes in 2018 and 2019 to investigate the effects of
96 water stratification and day-time sampling on Se distribution. To our knowledge, no studies or data
97 are available about Se in such aquatic systems. Our aim was thus both to contribute to
98 comprehensive data set of selenium occurrence and speciation in such pristine ecosystems and to
99 provide a unique “baseline” (Se background level) that allows to evaluate respective contribution
100 of headwaters and atmospheric inputs versus the impact of land uses in a specific watershed (Adour
101 river watershed on the French slope).

102

103 **2. Materials and Methods**

104 **2.1. Study site**

105 All the lakes considered in this study are located along a north-south transect of the Pyrenees (Fig.
106 1). Lakes can be divided into three sectors according to their location: Caunterets, Panticosa, and
107 Ossau/Ayous Valleys. Most of the studied lakes are located on granitic rocks (GR). The study area
108 lays on three different geological periods: Devonian, Permo-Triassic and Cretaceous (Zaharescu et
109 al., 2016; Gleizes et al., 1998; Zwart & De Sitter, 1979) which may impact Se bedrock
110 concentration. The geology of the Central Pyrenees has been described in detail elsewhere (Van

111 Lith, 1968; Catalan et al., 2006). In brief, granite (pDe-GR) originated during mountain formation
112 in the pre-Devonian geological age are found ubiquitously. Devonian sedimentary rocks (De-SR)
113 consisted of limestone, marble, shale and sandstone; Permo-Triassic bedrock (PT-SR) contains
114 sedimentary rocks in the form of conglomerates, sandstone and red or green shales; and, the most
115 recent formations, Cretaceous sedimentary rocks (Cr-SR) consist of limestone and sandstone.
116 Studied lakes are ice-covered during 5-6 months per year (Santoloria et al., 2015). Late spring
117 sampling (June/July) was thus scheduled to coincide with the thawing phase, to probe the effect of
118 recent ice melting and early lake productivity. Late summer-early fall sampling preceded holomixis
119 phase, to probe the effect of summer time stratification and whole biological production (both auto
120 and heterotrophic).

121

122 **2.2. Sampling lake waters and in-situ sample treatments**

123 A total of 71 subsurface water samples (0.2 – 0.5 m) were collected in 20 lakes during June/July
124 and October 2017–2019. Details of the sampling locations are summarized in Table 1. In brief, five
125 sampling campaigns were carried out in June/July 2017, October 2017, June 2018, October 2018
126 and June 2019. In the two first sampling campaigns of June and October 2017, three replicate
127 subsurface samples were collected in the upstream, central and downstream sectors of each lake to
128 examine intra-lake variabilities. Diurnal and depth profiles from lakes Gentau and Sabocos were
129 sampled in June 2018, October 2018 and June 2019. Azules diurnal and depth profiles were sampled
130 in June and October 2018, while Arratille profiles were sampled only in June 2018. The diurnal
131 cycle consisted in the acquisition of subsurface samples carried out every few hours from dawn to
132 sunset mainly to evaluate the effect of light in Se speciation.

133 For the Bachimaña reservoir and other lakes of difficult access, water samples were collected from
134 the shore, below the surface, using nitrile gloves to avoid contamination. For all other lakes, samples
135 were collected onboard an inflatable rubber boat using a non-metallic and PTFE coated sampler (5
136 L Go-Flo; General Oceanic). The sampler was operated with gloves for subsurface waters and fixed

137 on a Kevlar cable (scaled every meter from 0 to 40 m) for samples collected at depth. One litre of
138 the sample was immediately transferred into two 500 mL Teflon bottles with a pre-cleaned silicone
139 tubing, avoiding bubbles and head-space as for dissolved oxygen sampling, and stored in the dark
140 as fresh as possible for its subsequent purge and trap of the gaseous species.

141 Two aliquots either unfiltered, for bulk, or filtered, for dissolved, total Se analyses were collected
142 in 15 mL polypropylene tubes and acidified to 1% v/v HNO₃ (69%, trace metal grade). For total
143 dissolved Se (TDSe), samples were filtered (PVDF Sterivex filter units, 0.22µm, Millipore) prior to
144 their acidification. To study Se speciation, filtered water samples were stored without headspace in
145 50 mL Falcon tubes, as fresh as possible (4–10 °C) in the field in the dark. Once in the lab, all
146 samples were stored at ca. 4 °C in the dark until the analysis (less than two weeks). Additional
147 samples were collected and analyzed at the University of the Basque Country (Bilbao) for the main
148 anions, non-purgeable organic carbon (NPOC), and dissolved inorganic carbon (DIC) as previously
149 described elsewhere (Kortazar et al., 2020). Samples of depth profiles ($3 \leq n \leq 6$) and temporal
150 monitoring were treated following the same protocols detailed here and all procedures were
151 conducted trying to minimize their exposition to daylight.

152 The physicochemical parameters in water lakes were measured simultaneously to the sampling
153 using a multiparametric probe YSI EXO2 (temperature, conductivity, oxidation-reduction potential,
154 dissolved oxygen, turbidity). The pH was calculated according to Kortazar et al. (2020) with the
155 help of the CO2SYS software, this methodology providing a better precision and accuracy in
156 comparison with the pH measured by the probe.

157

158 **2.2. Total Se analysis**

159 Bulk total Se (TSe Bulk) was determined in unfiltered bulk samples (15 mL) after digestion with
160 300 µL HNO₃ (69%, tracepure grade) and 150 µL HCl (37%, tracepur grade) in sealed tubes and
161 incubated for 3 h at 90 °C in a hot block (DIGIPREP, SCP Science). The dilution factor was
162 corrected by weighting samples before and after the digestion.

163 Total Se concentrations were measured in bulk (TSe Bulk) and filtered (TDS_{Se}) samples with an
164 Agilent 7900x Series inductively coupled plasma mass spectrometer (ICP-MS) system (Agilent
165 Technologies, Tokyo, Japan) equipped with an octopole reaction cell, concentric nebulizer and a
166 Scott double pass spray chamber cooled to 2 °C. Argon-based polyatomic interferences were
167 reduced by using H₂ as cell gas at a flow rate of 5 mL min⁻¹ as previously optimised and validated
168 (Darrouzes et al., 2005). The parameters settings were as follow: Ar plasma gas flow, 15 L min⁻¹;
169 Ar auxiliary gas flow, 0.86 L min⁻¹; Ar nebulizer gas flow, 1–1.1 L min⁻¹; radio frequency (RF)
170 forward power, 1550 W. Acquisition parameters consisted of 10 replicates with 50 sweeps/replicate
171 and integration time of 2s per isotope; m/z monitored ratios were 77 and 78. External calibration
172 (range 5 – 100 ng L⁻¹ from single element Se standard 1000 mg L⁻¹ SCP Science) was performed.
173 Quality control used for quantification methodology was based on several repetitions (3 to 4 times)
174 of standard measurements during the day of analysis that led to calibration line slope relative
175 standard deviation <10%. Limits of quantification (based on ⁷⁸Se and according to IUPAC
176 recommandations) were between 0.7–2.1 ng L⁻¹. Typical analytical precision was <12% in the range
177 of measured concentrations of dozen ng Se L⁻¹ (relative standard deviation of the 10 replicates).

178

179 **2.3. Dissolved non-volatile Se speciation analysis**

180 Chromatographic separation was carried out with an Agilent 1200 HPLC pump hyphenated to ICP-
181 MS. Most samples were analyzed using a porous graphitic carbon stationary phase (Thermo
182 Hypercarb column 10 cm × 4.6 mm i.d) with a formic acid mobile phase (240 mmol L⁻¹, 1%
183 methanol and pH 2.4 adjusted with ammonia) delivered at 1 mL min⁻¹ flow rate (Dauthieu et al.,
184 2006). Chromatographic conditions allowed the separation of inorganic selenite and selenate and,
185 organic species (trimethylselenonium ion, methane seleninic acid, selenomethionine and
186 selenocystine) (Lanceleur et al., 2019; Be-eri-Shlevin et al., 2021). Standard addition was used for
187 quantification using daily diluted certified aqueous standards of 1 g L⁻¹ Se as selenite and 0.1 g L⁻¹

188 Se as selenate (Spectracer) (see example chromatogram Fig. SI 1). Quantification limits (LoQ) were
189 3.6 and 2.2 ng Se L⁻¹ for selenite (Se(IV)) and selenate (Se(VI)) respectively for 200 µl injected
190 volumes. Exceptionally, samples of October 2017 and part of those of October 2018 were analyzed
191 using the mixed-mode column OmniPac PAX-500 (Thermo, 25 cm x 4 mm i.d.) with a 20 mmol
192 L⁻¹ ammonium nitrate mobile phase containing 2% methanol at pH 8.0 (adjusted with ammonia),
193 delivered at 1 ml min⁻¹ flow rate. In this case, LoQ were 12 and 10 ng Se L⁻¹ for Se(IV) and Se(VI),
194 respectively, for 100 µl injected volumes. Duplicates of all samples were injected obtaining a
195 relative standard deviation below 10%, except for some samples close to the LoQ for which relative
196 standard deviation was up to 15%. Only selenate and eventually selenite were detected. The
197 operationally defined Se (-II, 0) fraction was then calculated by subtraction of species concentrations
198 from the total dissolved Se concentration.

199

200 **2.4. Dissolved gaseous Se speciation analysis**

201 Dissolved gaseous Se species were purged and trapped within the day of sampling in the field lab
202 set in a mountain hut. This allowed the pre-concentration of dissolved gaseous Se compounds and
203 the preservation of their speciation as described by Lanceleur et al. (2019). Samples (2x500 mL)
204 were purged with pure N₂ (500 mL min⁻¹) for 45 min. The resulting water vapour during the purge
205 was removed from the gas stream in a moisture trap maintained at -20 °C. The gas stream was then
206 carried through a volatile Se compounds trap (glass tube packed with Carbotrap sorbent). After the
207 purge, the glass columns were tightly closed with Teflon lined plugs and stored in the dark at 4 °C
208 in a sealed double PE bags until analysis. In the laboratory, samples were analyzed within a week
209 after sampling using a cryogenic GC-ICP/MS set-up. Samples were thermo-desorbed from
210 carbotraps at 250 °C for 2 minutes under He flow (100 mL min⁻¹). Samples were flushed and trapped
211 on the head of a Cryo GC column submerged in liquid N₂, prior to GC elution on Chromosorb
212 SP2100 (Amouroux et al., 1998). Quantification was obtained by external calibration. The limits of

213 quantification were between 1.3–3.8 pg Se L⁻¹ for DMSe and DMSeS and between 1.3–5.4 pg Se
214 L⁻¹ for DMDS₂Se. Thermodesorption efficiency was controlled by carrying out two consecutive
215 analyses of the same carbotrap column. TVSe content was determined as the sum of DMSe, DMSeS
216 and DMDS₂Se concentrations.

217

218 **2.5. Data processing and statistics**

219 In samples that did not contain quantifiable selenite and/or selenate (<LoQ), a value of one half their
220 quantification limit (with associated error of 100%) was used for Se (-II, 0) fraction calculation. The
221 limit of quantification of Se (-II, 0) fraction was calculated as the square root of the sum of the
222 squares of TDSe, Se(IV) and Se(VI) limits of quantification and, was in the range 4.5–15 ng Se L⁻¹.
223 For the statistical treatment, a value of one half the quantification limit (with associated error of
224 100%) was also assigned to the species (selenate, volatile species) and the Se (-II, 0) fraction
225 concentrations in samples that did not contain quantifiable levels.

226 All data sets were tested for normality using the Shapiro-Wilk test. The non-parametric pairwise
227 Wilcoxon test or Kruskal-Wallis followed by Conover-Iman tests for multiple, two-tailed,
228 comparisons were performed. Spearman coefficients were used for correlation analysis. These
229 analyses were carried out using using R Commander package (R Core Team, 2021, version 4.0.5)
230 and XLSTAT software (version 2021.2.1).

231

232 **3. Results and Discussion**

233 **3.1. Subsurface lake waters**

234 Detailed data for physicochemical characteristics and selenium concentrations are available in
235 Supporting information Tables SI 1 and SI 2. In June and October 2017 sampling campaigns, the
236 triplicate samples, collected in each lake to check intra-lake variabilities, exhibited no remarkable
237 differences between replicates (relative standard error of the mean was ≤13, 11 and, 9 %, respectively,
238 for bulk and dissolved total Se and Se(VI), data not shown). The diurnal cycle

239 measurements (June 2018, October 2018, and June 2019) also did not show significant variability
240 along the day for total Se, nor for speciation (data not shown). These results indicated primarily a
241 slow reactivity of Se compounds and Se turnover in the surface waters. We can thus assume that no
242 apparent bias was due to the sampling time of the day and that our measurements were representative
243 of sampled lakes in a given seasonal period.

244 *3.1.1 Physicochemical characteristics and selenium total concentrations*

245 Lake waters presented temperatures ranging from 2 to 19 °C (Table SI 1) and subsurface samples
246 were always oxygenated (>62% of oxygen saturation, data not shown). Conductivity ranged
247 between 4.5 and 130 $\mu\text{S cm}^{-1}$. The pH varied from slightly alkaline (7.91) to acidic values (4.87).
248 Oxidation-reduction potential (E_{H}) ranged from slightly reducing (36 mV) to oxidising (285 mV)
249 conditions. No significant differences were observed for conductivity, pH and E_{H} values between
250 June and October (Wilcoxon test, $p>0.05$). Some of the anions measured, such as fluoride (F^-),
251 bromide (Br^-), nitrites (NO_2^-) or phosphates (PO_4^{3-}), were frequently not detected (<LoQ, Table SI
252 1). Sulphate ($\leq 7.56 \text{ mg L}^{-1}$) was the main anion found in waters with a median value of 1.6 mg L^{-1}
253 ($n=71$). The highest sulphate concentrations were observed in Azul Superior (from 4.84 to 7.56 mg
254 L^{-1}) and Badète (from 2.96 to 5.43 mg L^{-1}) (Table SI 1). The median value found in October (1.89
255 mg L^{-1} , $n=27$) was slightly higher than in June (1.36 mg L^{-1} , $n=44$) but statistically not significantly
256 different (Wilcoxon test, $p=0.48$). Nitrate concentration was $\leq 1.16 \text{ mg L}^{-1}$ at all sites with highest
257 values found at Azul Superior (June 2019, 1.16 mg L^{-1}), Badète (June 2018, 1.13 mg L^{-1}) and
258 Bachimaña (June 2019, 1.08 mg L^{-1}) (Table SI 1). Nitrate median concentration of sub-surface
259 samples was 0.46 mg L^{-1} ($n=71$). The median value in June samples (0.56 mg L^{-1} , $n=44$) was almost
260 double that in October (0.29 mg L^{-1} , $n=27$) and significantly different (Wilcoxon test, $p=0.004$).
261 Dissolved organic matter ranged from 0.63 to 5.3 mg L^{-1} and median values were significantly
262 different between June (1.03 mg L^{-1} , $n=43$) and October (1.72 mg L^{-1} , $n=26$) (Wilcoxon test,
263 $p=0.0006$).

264 Total Se bulk concentrations ranged from 9 ± 1 to 82 ± 5 ng L⁻¹ while dissolved Se ranged between
265 7 ± 1 and 78 ± 7 ng L⁻¹ (Table SI 2). No significant differences were found for both concentrations
266 between June and October sampling campaigns (Wilcoxon test, $p > 0.05$) but bulk and filtered Se
267 concentrations were significantly different (Wilcoxon test, $p < 0.0001$). However most of the Se was
268 in the dissolved or colloidal phase in these lakes (Fig. SI 2). Measured dissolved Se concentrations
269 in Pyrenean lakes were similar to those reported in Norwegian lakes located on granitic gneisses
270 ($16\text{--}75$ ng L⁻¹) (Økelsrud et al., 2016). The highest total Se concentrations were found at Badète
271 (59.3 ng L⁻¹ on average) and Azul Superior lakes (57.8 ng L⁻¹ on average). Grouping lakes
272 according to underlying bedrock, the average dissolved Se concentration was 19 ± 6 ng L⁻¹ for lakes
273 located at the granitic core (pDe-GR) ($n=31$), 48 ± 18 ng L⁻¹ for Devonian sedimentary bedrocks
274 (De-SR) lakes ($n=21$), 12 ± 1 ng L⁻¹ at lakes located over Permo-Triassic sedimentary bedrock (PT-
275 SR) ($n=9$) and 14 ± 2 ng L⁻¹ for lake Sabocos ($n=3$) (Cretaceous prevailing bedrock, Cr-SR).
276 Selenium concentrations (bulk and dissolved) were thus significantly higher in De-SR lakes
277 compared to other bedrock lakes (Kruskall-Wallis and Conover-Iman tests, $p < 0.0001$) (Fig. 2A,
278 Table SI 2).

279 No relationship was found between total Se concentration and lake or catchment surface area. Lake
280 surface area demonstrating a linear relationship with water residence time (personal communication,
281 Lluís Camarero, CSIC, Spain), the lack of correlation between Se concentration and surface area
282 indicated that Se contents in lake waters were not drastically affected by water residence times.
283 Correlation analysis showed that total dissolved selenium concentrations were positively correlated
284 with those of sulphate, dissolved inorganic carbon and nitrates (Table SI 3, Fig. SI 3). Among them,
285 sulphate and dissolved inorganic carbon concentrations were also correlated (Table SI 3). The strong
286 link between total dissolved selenium concentrations and those of sulphate has been previously
287 reported in freshwaters with higher Se concentrations (> 100 ng L⁻¹, Ponton et al., 2020) and will
288 be further discussed in Paragraph 3.3. TDSe did not correlate with NPOC, meaning that Se was

289 weakly incorporated or bound to organic matter in such oligotrophic lakes. Very weak or no
290 correlation of TDSe with dissolved organic matter was previously reported in Siberian thaw lake
291 waters in spring time corresponding to frozen underlying peat preventing leaching of dissolved
292 organic matter and Se from peat layers (Pokrovsky et al., 2018).

293 *3.1.2 Selenium speciation*

294 Among non-volatile dissolved species, selenate was most often the only species detected with no
295 significant difference between June and October (Wilcoxon test, $p > 0.05$). The average values within
296 each lake group followed the same trend as for total dissolved Se (Fig. 2A), and were: 12 ± 7 (n=21),
297 42 ± 25 (n=15), 8 ± 2 (n=6) and 10.3 ± 0.4 (mean deviation, n=2) ng Se L⁻¹ for pDe-GR, De-SR,
298 PT-SR and Cr-SR groups, respectively for June sampling campaigns. After summer, selenate
299 concentration values were slightly lower: 10 ± 5 (n=14), 37 ± 17 (n=9), <10 (n=3) and 6 (n=1) ng
300 Se L⁻¹, respectively. Spring to late summer selenate concentrations decrease in meso-eutrophic
301 freshwater lake has been related to phytoplankton uptake, and was associated also to selenite
302 concentration decrease (Be'eri-Shlevin et al., 2021). Selenate, which is the stable form of Se in
303 oxic waters (Belzile et al., 2000), accounted for 64 ± 28 % of total dissolved Se, and both
304 concentrations were thus significantly different (Wilcoxon test, $p < 0.0001$). This was particularly
305 the case for pDe-GR, PT-SR and Cr-SR lake groups (Fig. SI 4) where other species may thus coexist
306 at extremely low concentration. This may be the case of selenite, which was typically undetected
307 ($<$ limit of quantification, in the range 3.6–12 ng Se L⁻¹) and the Se(-II, 0) fraction (LoQ in the range
308 4.5–15 ng Se L⁻¹). Pyrenean lakes are usually oligotrophic due to low contents of nutrient and
309 organic matter (Catalan et al., 2006), the presence of organic selenide was thus not expected as main
310 component of Se(-II, 0) fraction and no unknown selenium-containing compound could be
311 chromatographically detected (see example chromatogram Fig. SI 1). Considering E_H and pH values
312 observed in lake waters, elemental Se should be thermodynamically stable in most lakes (Seby et
313 al., 2001). Estimation of Se(-II, 0) fraction (Table SI 2) indicated the presence of such fraction in
314 both seasons in the range 4.7–17 ng L⁻¹ in subsurface waters at some lakes (Bachimaña Bajo,

315 Coanga and Baños de Panticosa pDe-GR lakes and, Ordicuso De-SR lake). Bibliographic data for
316 Se speciation in lakes is scarce, especially in remote lakes. Existing publications report selenate
317 concentrations in the range of 4–17 ng Se L⁻¹ in Finish lakes representing 6–12% of total Se, while
318 most of Se present was linked to organic matter (Wang et al., 1994; 1995). In Canadian lakes,
319 selenate was predominant (≥60%) in lakes where Se inputs occur principally in this form (Ponton
320 et al., 2013).

321 Total volatile Se ranged from 2.6 to 118 pg L⁻¹ with an exceptionally high value observed in June
322 2017 at Ordicuso (484 pg L⁻¹) without significant seasonal variation in each lake group (Fig. 2B
323 and Table SI 2). Concentrations showed similar wider range of values for pDe-GR and De-SR lakes
324 (4 to 118 pg L⁻¹ excluding Ordicuso in June 2017) compared to PT-SR and Cr-SR lakes (3 to 36 pg
325 L⁻¹). In pDe-GR lake groups, although not statistically different, average TVSe concentration in
326 June samples (66 ± 34 pg L⁻¹), just after the ice-cover melting, was higher compared to October
327 samples (41 ± 24 pg L⁻¹), which may indicate volatile species accumulation during ice capping
328 period. The main species found in all lakes was DMSe, representing in average 76 % of the TVSe
329 detected. Previous studies in remote arctic thermokarst ponds reported total volatile Se in the range
330 1–32 pg L⁻¹ including 96% of DMSe (Lanceleur et al., 2019). The low concentrations of TDSe and
331 sulphate in Pyrenean lakes seem to limit the production of dimethyldiselenide (DMDS₂) or dimethyl
332 selenyl sulfide (DMSeS) observed in other environments, such as meso-eutrophic lakes,
333 minerotrophic peatlands or estuarine and marine systems (Amouroux & Donard, 1997; Tessier et
334 al., 2002; Vriens et al., 2014; Be'eri-Shlevin et al., 2021). For TVSe, no clear trend was observed
335 with any of the measured parameters in water samples (Spearman coefficient $|\rho| < 0.3$), most
336 probably because of the extremely low concentration of gaseous Se species in most lakes. The
337 limited nutrient supplement and further biological activity of Pyrenean lakes seem to be the main
338 limit for volatile Se production. In addition, the short lifetime and rapid degradation of volatile Se

339 compounds in the photic zone could make it difficult to establish correlation trends (Mason et al.,
340 2018).

341

342 **3.2. Profiles of selenium and dissolved species in lake waters**

343 During the sampling campaigns of 2018 and 2019, depth profiles were carried out in four lakes:
344 Arratille and Azul Superior (De-SR lakes), Gentau (PT-SR lake) and Sabocos (Cr-SR lake).
345 Detailed data are available in Table SI 4.

346 *3.2.1 Shallow lakes (maximum depth < 15 m)*

347 Arratille and Azul Superior lakes belong respectively to Cauteret and Panticosa areas of similar
348 geological structure (mostly granitic and, limestone, marble, shale and sandstone sedimentary rocks)
349 but respectively located in the northern and southern slope. Oxygen and temperature depth profiles
350 demonstrate that Arratille exhibited a small oxygen depletion only at the bottom water sample in
351 June, while Azul Superior, with a maximum depth of around 8 m, had a well-mixed water column
352 (Fig. SI 5). In both lakes, total Se concentration did not vary significantly and the main species was
353 selenate with less than 6 and 7% variation, respectively, along the water column. Selenite and Se(-
354 II, 0) fraction were both below the LoQ. The profile of volatile species was different from those of
355 non-volatile selenium species. In June, the concentration of DMSe was increased at half-depth in
356 both lakes. This concentration peak was more visible in Aratille lake where DMSe concentration
357 stayed constant to the bottom. In Azul Superior lake, DMSe profile appeared to be similar to the
358 one of nitrate in June while it was opposite in October where DMSe concentrations were clearly
359 lower. The formation of DMSe has been shown to occur with similar production rates from selenate
360 or selenite in bottom pond waters (i.e oxygen depleted) (Lanceleur et al., 2019). Although, DMSe
361 production in the photic zone could not be excluded, it should be of lesser extent in such oligotrophic
362 lakes if compared to deeper microbial processes in bottom waters and sediments (Amouroux &
363 Donard, 1997; Lanceleur et al., 2019). Minor volatile algal production was also previously observed

364 comparing marine algae, bacteria and mixed bacteria-algae culture experiments (Luxem et al.,
365 2017). Observed difference between sub-surface and bottom volatile Se concentrations could thus
366 be indicative of its formation and accumulation at depth during winter-early spring (i.e ice-covered
367 period promoting reducing conditions), upward diffusion and volatilization and/or photodegradation
368 in the surface layer during summer period.

369 *3.2.2 Deep lakes*

370 Gentau and Sabocos lakes are located at lower elevation and belong respectively to Ayous and
371 Panticosa areas where agropastoralism activities can represent a source of organic matter and
372 nutrients (Kortazar et al., 2020). Both lakes are affected by thermal and oxygen stratification and
373 during 2018 samplings, the deepening of thermocline and oxycline was observed in both lakes from
374 late spring to autumn (Fig. 3). Hypoxic conditions ($\text{DO} < 4 \text{ mg L}^{-1}$) were maintained in the deepest
375 5 m. Nitrate and sulphate concentrations in Sabocos were a little lower than the mean values (≈ 1
376 and 4 mg L^{-1} , respectively) measured in 2012 (Santoloria et al., 2015) and, higher compared to
377 Gentau. This was particularly the case for sulphate concentration that was extremely low ($\leq 0.7 \text{ mg}$
378 L^{-1}) in Gentau. After summer, a nitrate depletion was observed in the photic zone (4–8 m depth in
379 Sabocos, up to 12 m depth in Gentau). Conductivity values were also higher in Sabocos due to its
380 calcareous basin (Santoloria et al., 2015; Kortazar et al., 2020). An increase of conductivity values
381 from the surface to the bottom was observed in Gentau. With the exception of two deviant values
382 observed in Gentau (abnormally high values for TDS_{Se} at 0.5 m depth in June 18 and, TSe bulk at 18
383 m depth in June 19), total Se concentrations did not vary significantly along the water column ($<$
384 14% variation) as previously observed in shallower lakes.

385 Selenite concentration was in both lake profiles below the LoQ and, selenate demonstrated opposite
386 profiles. Selenate was not detected in October profile of Gentau lake which may be related to the
387 high LoQ of 10 ng L^{-1} of the chromatographic conditions used. However, reproducible profiles were
388 observed in June where selenate concentration decreased with increasing depth, resulting in

389 quantifiable Se(-II, 0) fraction in hypoxic deepest 5 m. These data suggested either abiotic and/or
390 microbial reduction of selenate or its slower re-oxidation in bottom waters of Gentau lake as
391 observed in other lentic systems (Lanceleur et al., 2019; Be'eri-Shlevin et al., 2021). In Sabocos,
392 however, selenate proportion in June profiles did not vary significantly along the water column (67
393 ± 7 and 90 ± 9 % of total dissolved Se in 2018 and 2019, respectively). The lower proportion of
394 selenate in June 2018 resulted in occasional quantifiable Se(-II, 0) fraction at values marginally
395 above the LoQ. In October, despite an oxygen saturation above 70%, Se(VI) depletion was observed
396 in the photic zone as was the case for nitrate. These data could indicate biological uptake of selenate
397 as previously suggested elsewhere (Harrison et al., 1988; Duan et al., 2010; Be'eri-Shlevin et al.,
398 2021). Observed differences between selenate proportion in oxygen-depleted bottom waters of both
399 lakes may be related to the fact that Gentau lake receives mineral inputs, mainly iron (Kortazar et
400 al., 2020), which may promote the existence of Fe-selenides or colloidal elemental selenium at the
401 sediment-water interface (Fernández-Martínez & Charlet, 2009).

402 The total volatile Se concentration sharply peaked in both bottom waters in June 2018 at values that
403 were not measured afterward. This increase mainly resulted from DMDSe contribution (43 to 76%
404 of TVSe). In comparison with shallower lakes, DMSeS contribution to total volatile selenium was
405 non negligible in the photic zone of Gentau and Sabocos lakes. In addition to enhanced production
406 of DMSe, Lanceleur et al. (2019) have reported that DMDSe was formed in bottom pond waters
407 enriched with local biofilm with a production rate approximately 100 times lower compared to the
408 one of DMSe. The significant proportion of DMDSe with respect to DMSe observed in-depth of
409 these oligotrophic aquatic systems may thus be related to their respective volatility (Karlson et al.,
410 1994) and dark conditions favourable to formation and stability of diselenide bond (Amouroux et
411 al., 2000). The presence of DMSeS in shallower waters suggested its possible formation from
412 DMDSe and sulphur-containing volatile compounds (Chasteen, 1993; Swearingen et al., 2006).
413 Likewise previous observations in shallow lakes, enhanced concentration levels of TVSe were

414 observed in springtime after the ice coverage period preserving such compounds from volatilization
415 and photodegradation that seem to control TVSe in sub-surface waters.

416
417 **3.3. Sources of Selenium in alpine lake waters: bedrocks leaching/erosion versus wet**
418 **depositions inputs**

419 Because most of the investigated lakes have a limited watershed size (Table 1), main Se inputs are
420 suggested to be related to the elemental composition of the prevailing bedrocks through dissolution
421 and erosion pathways, and direct atmospheric inputs. Among major anions and considering all lakes
422 and sampling dates, correlation analysis showed a very strong positive relationship between total
423 dissolved Se and sulphate (Fig. SI 3). Sulphate's concentration mainly depends on the geological
424 substrate while the intensive study of Camarero et al. (2009) in European mountain lakes revealed
425 that lakes having sulphate concentration $< 2.3 \text{ mg L}^{-1}$ could be considered as mainly controlled by
426 S atmospheric inputs. During our recent field sampling program, a similar threshold sulphate
427 concentration value of 0.7 mg L^{-1} has been observed and proposed to distinguish between
428 atmospheric or geological sources in lakes of studied zone (Duval, 2020). This value agrees with
429 the annual sulphate concentration of 0.67 mg L^{-1} reported by Suess et al. (2019) at the Pic du Midi
430 de Bigorre high altitude station. In the western Pyrenees region, shales formed during the Devonian
431 geological period are the main source of SO_4^{2-} (Van Lith, 1968). Lakes presenting the highest
432 sulphate and selenium concentrations are those located over sedimentary rocks containing shales
433 (De-SR lakes group). Their SO_4^{2-} concentration is higher than concentrations found in rainwaters
434 (Roulier et al., 2021; Suess et al., 2019) that may thus, similarly to S, indicate selenium geological
435 supply as selenium concentrations are typically higher in shales than in granite, limestone or
436 sandstone (Plant et al., 2014; Fernandez-Martinez & Charlet, 2009).

437 To evaluate the contribution of geogenic versus atmospheric Se sources, the molar ratio of Se/S
438 measured in lake waters in this study was compared with Se/S ratios calculated for different rock
439 types using a standard database (Reimann & De Caritat, 1998) as no data was available for local

440 bedrocks. Molar Se/S ratios measured in lake waters were also compared with Se/S ratios in wet
441 deposition at the Pic du Midi de Bigorre (calculated from Suess et al., 2019) and Adour watershed
442 (calculated from Roulier et al., 2021). The molar ratios Se/S in Pyrenean lakes were in the range
443 $(0.7\text{--}6.8) \cdot 10^{-5}$ (average values per lake groups in Table 2) and do not allow to provide clear
444 discrimination of potential major sources of Se in those lakes. However, considering the threshold
445 value of sulphate concentration in rain waters ($0.7 \text{ mg L}^{-1}=7.5 \text{ } \mu\text{mol L}^{-1}$) below which wet
446 depositions is the main source of S (Duval, 2020; Camarero et al., 2009), a strong direct relationship
447 was obtained between Se/S ratio and sulphate concentration for half of the studied lakes belonging
448 to pDe-GR and PT-SR groups (Spearman coefficient $\rho=-0.75$, $p<0.0001$, Fig. 4). For the other half
449 of lakes, including all De-SR group lakes, for which sulphate concentration was higher than 7.5
450 $\mu\text{mol L}^{-1}$ and controlled by geological substrate, only a weak correlation was obtained (Spearman
451 coefficient $\rho=-0.34$, $p<0.05$, Fig. 4).

452 An estimation of Se input from wet deposition was calculated using the average annual Se flux in
453 Adour watershed located downstream the northern section of the west Pyrenees ($50 \pm 20 \text{ } \mu\text{g Se m}^{-2}$
454 yr^{-1} , Roulier et al., 2021). The estimation considered the surface area and volume of each lake to
455 compare the input of Se from wet deposition with the stock of Se at each lake. The lake volume and
456 surface area data were taken from Duval (2020) and the results are presented in Table SI 5. We
457 found that Se provided from annual wet deposition could account for 12 to 100% (median value of
458 39%) of Se stored in lake water at a specific time of the year. Thus, wet depositions through rain or
459 snow events and during the snowmelt period is a non-negligible source of selenium in such alpine
460 Pyrenean lakes, which can be counterbalanced by a Se-enriched geological substrate.

461 The emission of volatile Se compounds could then contribute to lower the extent of Se in surface
462 lake waters, mainly in the form of DMSe. We used our measurements to estimate Se gaseous output
463 from lake water to the atmosphere, but as they corresponded only to two months of the year the
464 variability of TVSe concentrations was not well constrained. However, due to the low TVSe

465 concentrations and low seasonal variability observed before and after summer, together with the
466 low surface area of most lakes (Table 1), low volatilization fluxes are expected. A rough estimation
467 of Se emission was carried out similarly to Lanceleur et al. (2019) using the model of Cole & Caraco
468 (1998) developed for small lakes considering 3 m s^{-1} as maximum average wind speed, as recorded
469 in those sheltered lake systems (ca. $1\text{--}3 \text{ m s}^{-1}$, from Duval et al., in preparation). We assumed that
470 atmospheric exchange occurred for approximately 6 months, while during late fall to late spring the
471 ice cover prevented volatilization. The annual Se emission (Table SI 5) was compared with the
472 estimated total Se stock in the corresponding lake. These values ($3.8\text{--}31 \text{ ng Se day}^{-1} \text{ m}^{-2}$) were
473 similar to the daily emission reported at thermokarst ponds ($1\text{--}97 \text{ ng Se day}^{-1} \text{ m}^{-2}$) (Lanceleur et
474 al., 2019). The results indicated that annual Se emission from lakes to the atmosphere
475 approximately represents 0.2 to 6.4% of the Se stock in lakes at a specific time of the year (median
476 value of 3%), remaining thus a minor pathway for Se removal from lake waters.

477

478 **4. Conclusions**

479 Pyrenean lakes showed similar bulk and dissolved total Se concentrations, in the low range of a
480 dozen to a few dozen of ng Se L^{-1} . Total dissolved Se varied between $7\text{--}78 \text{ ng L}^{-1}$ and, on average,
481 64% was in the form of selenate in subsurface samples. The extremely low Se concentration in
482 alpine lakes limited the detection of other Se species such as selenite. Volatile Se speciation was
483 mainly limited to the production of DMSe and volatilization appeared to not represent an important
484 removal pathway for Se in Pyrenean lakes.

485 Since no significant seasonal variations (June versus October) was observed neither for total Se
486 concentrations nor its species in subsurface waters, it is suggested that Se concentrations in lake
487 waters are regulated by downstream export and possible scavenging. The proportion of Se (-II, 0)
488 fraction was for example increased in-depth of deep lakes, such as Gentau lake, together with
489 oxygen depletion probably because of microbial reduction. Further studies are required to ascertain

490 how this process promotes Se scavenging, precipitation and further sedimentation in those lakes, as
491 shown in thermokarst artic ponds (Lanceleur et al., 2019).

492 The values of molar ratios Se/S in Pyrenean lakes overlap those for comparable bedrock types and
493 wet depositions. Our interpretation suggests that atmospheric inputs represent an important source
494 of Se to remote lakes when bedrock geochemistry is depleted with Se and sulphate. This study
495 provides a unique set of data that allows to better assess Se biogeochemistry in lakes and watershed
496 mainly affected by global emissions and long-range transport and atmospheric depositions.

497

498 **Conflicts of interest**

499 There are no conflicts of interests to declare.

500

501 **Acknowledgements**

502 This work is a contribution to the REPLIM-OPCC project and has been partially supported (65%)
503 by the FEDER funds through the INTERREG V-A Spain-France-Andorra (POCTEFA 2014-2020)
504 (REPLIM project, ref. EFA056/15). A. Romero–Rama thanks the UPPA and IPREM for her PhD
505 grant (Sciences Doctoral School, UPPA). The contributions of the Aquitaine Region
506 (AQUITRACES project n° 20131206001-13010973) and ANR IA RSNR (AMORAD project
507 n°ANR-11-RSNR-0002) for equipment funding are also acknowledged. Ainara Gredilla and Olaia
508 Liñero Campo (University of the Basque Country) are acknowledged for the analysis of anions.

509

510 **5. Bibliography**

- 511 D. Amouroux, O.F.X. Donard, Evasion of selenium to the atmosphere via biomethylation processes
512 in the Gironde estuary, France, *Mar. Chem.*, 1997, **58**, 173–188.
- 513 D. Amouroux, E. Tessier, C. Pécheyrán, O.F.X. Donard, Sampling and probing volatile metal(loid)
514 species in natural waters by in-situ purge and cryogenic trapping followed by gas
515 chromatography and inductively coupled plasma mass spectrometry (P-CT-GC-ICP/MS),
516 *Anal. Chim. Acta.*, 1998, **377**, 241–254.
- 517 D. Amouroux, C. Pécheyrán, O.F.X. Donard, Formation of volatile selenium species in synthetic
518 seawater under light and dark experimental conditions, *App. Organomet. Chem.*, 2000, **14**, 236-
519 244.
- 520 Y. Be'eri-Shlevin, M. Bueno, E. Tessier, A. Romero-Rama, A. Sukenik, T. Zohary, D. Amouroux,
521 Biogeochemistry of selenium compounds in the water column of warm monomictic Lake
522 Kinneret, *Biogeochemistry*, <https://doi.org/10.1007/s10533-021-00877-6>.
- 523 N. Belzile, Y.W. Chen, R. Xu, Early diagenetic behaviour of selenium in freshwater sediments, *App.*
524 *Geochem.*, 2000, **15**, 1439–1454.
- 525 T. Blazina, A. Läderach, G.D. Jones, H. Sodemann, H. Wernli, J.W. Kirchner, L.H.E.E. Winkel,
526 Marine Primary Productivity as a Potential Indirect Source of Selenium and Other Trace
527 Elements in Atmospheric Deposition, *Environ. Sci. Technol.*, 2017, **51**, 108–118.
- 528 J.E. Brandt, E.S. Bernhardt, G.S. Dwyer, R.T. Di Giulio, Selenium Ecotoxicology in Freshwater
529 Lakes Receiving Coal Combustion Residual Effluents: A North Carolina Example, 2017,
530 *Environ. Sci. Technol.*, **51**, 2418–2426.
- 531 L. Camarero, M. Rogora, R. Mosello, N.J. Anderson, A. Barbieri, I. Botev, A.F. Lotter,
532 Regionalisation of chemical variability in European mountain lakes, *Freshw. Biol.*, 2009, **54**,
533 2452–2469.
- 534 J. Catalan, M. Ventura, A. Brancelj, I. Granados, H. Thies, U. Nickus, E. Stuchlík, Seasonal
535 ecosystem variability in remote mountain lakes: implications for detecting climatic signals in
536 sediment records, *J. Paleolimnol.*, 2002, **28**, 25–46.
- 537 J. Catalan, L. Camarero, M. Felip, S. Pla, M. Ventura, T. Buchaca, E.O. Casamayor, High mountain
538 lakes: extreme habitats and witnesses of environmental changes, *Limnetica*, 2006, **25**, 551–
539 584.
- 540 J.J. Cole, N.F. Caraco, Atmospheric exchange of carbon dioxide in a low-wind oligotrophic lake
541 measured by the addition of SF₆, *Limnol. Oceanogr.*, 1998, **43**, 647–656.
- 542 J.E. Conde, M. Sanz Alaejos, Selenium Concentrations in Natural and Environmental Waters,
543 *Chem. Rev.*, 1997, **97**, 1979–2004.

544 T.G. Chasteen, Confusion between dimethyl selenenyl sulfide and dimethyl selenone released by
545 bacteria, *App. Organomet. Chem.*, 1993, **7**, 335-342.

546 J. Darrouzes, M. Bueno, G. Lespes, M. Potin-Gautier, Operational optimisation of ICP-octopole
547 collision/reaction cell-MS for applications to ultratrace selenium total and speciation
548 determination, *J. Anal. At. Spectrom.*, **20**, 2005, 88–94.

549 M. Dauthieu, M. Bueno, J. Darrouzes, N. Gilon, M. Potin-Gautier, Evaluation of porous graphitic
550 carbon stationary phase for simultaneous preconcentration and separation of organic and
551 inorganic selenium species in “clean” water systems, *J. Chromatogr. A*, 2006, **1114**, 34–39.

552 L. Duan, J. Song, X. Li, H. Yuan, S. Xu, Distribution of selenium and its relationship to the eco-
553 environment in Bohai Bay seawater, *Mar. Chem.*, 2010, **121**, 87–99.

554 B. Duval, Ecodynamics of metals and metalloids in Pyrenean lakes in relation to climate change
555 and anthropogenic pressure, Ph.D. Thesis, Universite de Pau et des Pays de l’Adour and Euskal
556 Herriko Unibertsitatea, 2020.

557 A. Fernández-Martínez, L. Charlet, Selenium environmental cycling and bioavailability : a
558 structural chemist point of view, *Rev. Environ. Sci., Biotechnol.*, 2009, **8**, 81–110.

559 S. Gascoin, O. Hagolle, M. Huc, L. Jarlan, J.-F. Dejoux, C. Szczypta, R. Sánchez, A snow cover
560 climatology for the Pyrenees from MODIS snow products, *Hydrol. Earth Syst. Sci.*, 2015, **19**,
561 2337–2351.

562 G. Gleizes, D. Leblanc, V. Santana, P. Olivier, J.L. Bouchez, Sigmoidal structures featuring dextral
563 shear during emplacement of the Hercynian granite complex of Cauterets–Panticosa
564 (Pyrenees), *J. Struct. Geol.*, 1998, **20**, 1229–1245.

565 P.J. Harrison, P.W. Yu, P.A. Thompson, N.M. Price, D.J. Phillips, Survey of selenium requirements
566 in marine phytoplankton, *Mar. Ecol. Prog. Ser.*, 1988, **47**, 89–96.

567 U. Karlson, W.T. Frankenberger Jr., W.F. Spencer, Physicochemical properties of dimethyl selenide
568 and dimethyl diselenide, *J. Chem. Eng. Data*, 1994, **39**, 608–610.

569 L. Kortazar, B. Duval, O. Liñero, O. Olamendi, A. Angulo, D. Amouroux, A. de Diego, L.A.
570 Fernandez, Accurate determination of the total alkalinity and the CO₂ system parameters in
571 high-altitude lakes from the Western Pyrenees (France-Spain), *Microchem. J.*, 2020, **152**,
572 104345.

573 L. Lancelleur, E. Tessier, M. Bueno, R. Pienitz, F. Bouchard, C. Cloquet, D. Amouroux, Cycling
574 and atmospheric exchanges of selenium in Canadian subarctic thermokarst ponds,
575 *Biogeochemistry*, 2019, **145**, 193–211.

576 K. Lindström, *Peridinium cinctum* bioassays of Se in Lake Erken, *Arch. Hydrobiol.*, 1980, **89**, pp.
577 110–117.

- 578 J.L. López-Moreno, S.M. Vicente-Serrano, Atmospheric circulation influence on the interannual
579 variability of snow pack in the Spanish Pyrenees during the second half of the 20th century,
580 *Hydrol. Res.*, 2007, **38**, 33–44.
- 581
- 582 K.E. Luxem, B. Vriens, R. Behra, L.H.E. Winkel, Studying selenium and sulfur volatilisation by
583 marine algae *Emiliana huxleyi* and *Thalassiosira oceanica* in culture, *Environ. Chem.*, 2017,
584 **14**, 199–206.
- 585 R.P. Mason, A.L. Soerensen, B.P. diMento, P.H. Balcom, The global marine selenium cycle:
586 insights from measurements and modeling, *Global Biogeochem. Cycles*, 2018, **32**, 1720–1737.
- 587 Y.V. Nancharaiah, P.N.L. Lens, Ecology and biotechnology of selenium-respiring bacteria,
588 *Microbiol. Mol. Biol. Rev.*, 2015, **79**, 61–80.
- 589 A. Nishri, I.B. Brenner, G.E.M. Hall, H.E. Taylor, Temporal variations in dissolved selenium in
590 Lake Kinneret (Israel), *Aquat. Sci.*, 1999, **61**, 215–233.
- 591 A. Økelsrud, E. Lydersen, E. Fjeld, Biomagnification of mercury and selenium in two lakes in
592 southern Norway, *Sci. Total Environ.*, 2016, **566**, 596–607.
- 593 Y. P. Pan, Y.S. Wang, Atmospheric wet and dry deposition of trace elements at 10 sites in Northern
594 China, *Atmos. Chem. Phys.*, 2015, **15**, 951–972.
- 595 J.A. Plant, J. Bone, N. Voulvoulis, D.G. Kinniburgh, P.L. Smedley, F.M. Fordyce, B. Klinck,
596 Arsenic and Selenium, in *Treatise on Geochemistry*, 2nd ed., ed. K.H. Turekian and H.D.
597 Holland, Elsevier, Oxford, 2014, pp. 13–57.
- 598 O.S. Pokrovsky, M. Bueno, R.M. Manasypov, L.S. Shirokova, J. Karlsson, D. Amouroux, Dissolved
599 organic matter controls on seasonal and spatial selenium concentration variability in thaw lakes
600 across a permafrost gradient, *Environ. Sci. Technol.*, 2018, **52**, 10254–10262.
- 601 D.E. Ponton, L. Hare, Relating selenium concentrations in a planktivore to selenium speciation in
602 lakewater, *Environ. Pollut.*, 2013, **176**, 254–260.
- 603 D.E. Ponton, S.D. Graves, C. Fortin, D. Janz, M. Amyot, M. Schiavon, Selenium interactions with
604 algae : chemical processes at biological uptake sites, bioaccumulation, and intracellular
605 metabolism, *Plants*, 2020, **9**, 528.
- 606 C. Reimann, P. De Caritat, *Chemical elements in the environment: factsheets for the geochemist
607 and environmental scientist*, Springer Berlin Heidelberg, 1998.
- 608 H. Robberecht, R. Van Grieken, M. Van Sprundel, D. Vanden Berghe, H. Deelstra, Selenium in
609 environmental and drinking waters of Belgium, *Sci. Total Environ.*, 1983, **26**, 163–172.
- 610 M. Roulier, M. Bueno, F. Coppin, M. Nicolas, Y. Thiry, F. Rigal, I. Le Hecho, F. Pannier,
611 Atmospheric iodine, selenium and caesium wet depositions in France: I. Spatial and seasonal

612 variations, *Chemosphere*, 2021, **273**, <https://doi-org./10.1016/j.chemosphere.2020.128971>.

613 Z. Santoloria, T. Arruebo, J.S. Urieta, F.J. Lanaja, A. Pardo, J. Mantesanz, C. Rodriguez-Casals,
614 Hydrochemistry dynamics in remote mountain lakes and its relation to catchment and
615 atmospheric features: the case study of Sabocos Tarn, Pyrenees, *Environ. Sci. Pollut. Res.*,
616 2015, **22**, 231–247.

617 F. Seby, M. Potin-Gautier, E. Giffaut, G. Borge, O.F.X. Donard, A critical review of thermodynamic
618 data for selenium species at 25°C, *Chem. Geol.*, 2001, **171**, 173–194.

619 V.K. Sharma, T.J. McDonald, M. Sohn, G.A.K. Anquandah, M. Pettine, R. Zboril, Biogeochemistry
620 of selenium. A review, *Environ. Chem. Lett.*, 2014, **13**, 49–58.

621 D.B.D. Simmons, D. Wallschläger, A critical review of the biogeochemistry and ecotoxicology of
622 selenium in lotic and lentic environments, *Environ. Toxicol. Chem.*, 2005, **24**, 1331–1343.

623 E. Suess, F. Aemisegger, J.E. Sonke, M. Sprenger, H. Wernli, L.H.E. Winkel, Marine versus
624 Continental Sources of Iodine and Selenium in Rainfall at Two European High-Altitude
625 Locations, *Environ. Sci. Technol.*, 2019, **53**, 1905–1917.

626 J.W. Swearingen Jr., D.P. Frankel, D.E. Fuentes, C.P. Saavedra, C.C. Vásquez, T.G. Chasteen,
627 Identification of biogenic dimethyl selenodisulfide in the headspace gases above genetically
628 modified *Escherichia coli*, *Anal. Biochem.*, 2006, **348**, 115–122.

629 D. Tanzer, K.G. Heumann, Determination of Dissolved Selenium Species in Environmental Water
630 Samples Using Isotope Dilution Mass Spectrometry, *Anal. Chem.*, 1991, **63**, 1984–1989.

631 E. Tessier, D. Amouroux, O.F.X. Donard, Biogenic volatilization of trace elements from European
632 estuaries, in *Biogeochemistry of Environmentally Important Trace Elements*, ACS Symposium
633 Series, American Chemical Society, 2002, **835**, 151–165.

634 J. G. J. Van Lith, Geology of the Spanish part of the Gavarnie Nappe (Pyrenees) and its underlying
635 sediments near Bielsa (Province of Huesca), *Geologica Ultraiectina*, Utrecht University, 1968,
636 Vol. 10.

637 B. Vriens, A.A. Ammann, H. Hagendorfer, M. Lenz, M. Berg, L.H.E. Winkel, Quantification of
638 methylated selenium, sulfur, and arsenic in the environment, *PLoS ONE*, 2014, **9**.
639 <https://doi.org/10.1371/journal.pone.0102906>

640 D. Wang, G. Alfthan, A. Aro, Determination of Total Selenium and Dissolved Selenium Species in
641 Natural Waters by Fluorometry, *Environ. Sci. Technol.*, 1994, **28**, 383–387.

642 D. Wang, G. Alfthan, A. Aro, A. Mäkelä, S. Knuutila, T. Hammar, The impact of selenium
643 supplemented fertilization on selenium in lake ecosystems in Finland, *Agric. Ecosyst. Environ.*,
644 1995, **54**, 137–148.

645

- 646 D.G. Zaharescu, P.S. Hooda, C.I. Burghilea, A. Palanca-Soler, A multiscale framework for
647 deconstructing the ecosystem physical template of high-altitude lakes, *Ecosystems*, 2016, **19**,
648 1064–1079.
- 649 H.J. Zwart, L.U. De Sitter, The geology of the Central Pyrenees. *Leidse Geologische Mededelingen*,
650 1979, **50**, 1–74.