



HAL
open science

Characteristic features of toxic metal content in hair samples of foreign students at RUDN University from different geographic regions

Anatoly Kirichuk, Anatoly Skalny, Aleksey Ya. Chizhov, Alexander Tskhovrebov, Dirk Schaumlöffel, Andreii Kritchenkov

► To cite this version:

Anatoly Kirichuk, Anatoly Skalny, Aleksey Ya. Chizhov, Alexander Tskhovrebov, Dirk Schaumlöffel, et al.. Characteristic features of toxic metal content in hair samples of foreign students at RUDN University from different geographic regions. *Journal of Trace Elements in Medicine and Biology*, 2023, 80, pp.127303. 10.1016/j.jtemb.2023.127303 . hal-04224486

HAL Id: hal-04224486

<https://univ-pau.hal.science/hal-04224486>

Submitted on 2 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

**Characteristic Features of Toxic Metal Content in Hair Samples of Foreign Students at
RUDN University from different geographic regions**

*Anatoly A. Kirichuk^a, Anatoly V. Skalny^{a,b}, Aleksey Ya. Chizhov^a, Alexander G. Tskhovrebov^a,
Dirk Schaumlöffel^{a,c}, and Andreii S. Kritchenkov^{a,d,*}*

*^a Peoples' Friendship University of Russia (RUDN University), 6 Miklukho-Maklaya Street,
Moscow, 117198, Russian Federation*

^b Sechenov University, Moscow, 119435, Russian Federation

*^c CNRS, Université de Pau et des Pays de l'Adour, E2S UPPA, Institut des Sciences Analytiques
et de Physico-Chimie pour l'Environnement et les Matériaux (IPREM), UMR 5254, 64000 Pau,
France*

*^d Institute of Technical Acoustics NAS of Belarus, Ludnikova Prosp. 13, Vitebsk, 210009, Republic
of Belarus*

** Corresponding authors at Peoples' Friendship University of Russia (RUDN University),
Miklukho-Maklaya str. 6, Moscow, 117198, Russian Federation.*

E-mail address: kritchenkov-as@rudn.ru (Andreii S. Kritchenkov)

Tel.: +79819365895

Fax: +74959540336

Abstract

1 *Background:* This study explores the regional variations in toxic metal accumulation among
2 RUDN University students from various global regions.
3

4 *Methods:* This comparative analysis examined hair samples from students hailing from different
5 regions, including Russia, Asia, the Middle East, Africa, and Latin America. The concentrations
6 of Aluminium (Al), Arsenic (As), Cadmium (Cd), Mercury (Hg), Lead (Pb), and Tin (Sn) were
7 measured in the hair samples. The data was then evaluated using regression models to assess the
8 link between the region of residence and toxic metal content in the hair.
9

10 *Results:* The analysis indicated significant regional variations in the levels of toxic metals in the
11 students' hair. The highest content of Al, Cd, and Pb was observed in students from Africa (13.542,
12 0.028, 0.794 $\mu\text{g/g}$) and Latin America (9.947, 0.025, 0.435 $\mu\text{g/g}$). Arsenic levels in students from
13 all regions exceeded that of Russian students by over two-fold. No substantial group differences
14 were found in the Sn content. The regression models suggested that residing in Asia, Africa, and
15 Latin America was a predictor of high Hg levels in hair (0.130, 0.096, 0.227 $\mu\text{g/g}$). Living in Africa
16 was significantly associated with higher Pb levels (0.794 $\mu\text{g/g}$), and living in Latin America was
17 close to significantly associated with the Cd level in the hair (0.025 $\mu\text{g/g}$).
18

19 *Conclusion:* This study confirmed an increased accumulation of toxic metals, especially Hg, Cd,
20 and Pb, in students primarily from Latin America and Africa. The findings highlighted the
21 importance of understanding the regional variations in toxic metal accumulation to address
22 associated health risks and the potential impact on students' well-being and academic performance.
23 These insights may guide the development of targeted interventions to reduce exposure to toxic
24 metals in students from various regions around the world.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

The intensive development of mining and heavy industry, road and air transport, the burning of coal and other materials over the past century has inevitably led to environmental pollution with toxic metals [1-7]. Toxic metals, including mercury (Hg), lead (Pb), cadmium (Cd), aluminum (Al), arsenic (As), and tin (Sn), have been widely recognized as harmful substances that can accumulate in the human body and cause a variety of health issues, such as neurotoxicity, nephrotoxicity, and cardiovascular diseases [8-19]. In many instances, exposure to toxic metals results in severe behavioral disorders, decreased intelligence, and learning ability [20-22]. The accumulation of these toxic elements in humans varies significantly across different regions of the world, which is primarily due to the differences in environmental pollution levels and exposure pathways in these areas [23, 24].

To assess the levels of toxic elements in the human body, various analytical methods involving urine, blood, and hair samples have been employed [25, 26]. Among these methods, hair analysis offers several advantages. Hair samples are non-invasive, easier to collect, and require minimal processing and storage conditions compared to blood and urine samples [27]. Moreover, while blood analysis provides information on the current state of toxic element exposure, hair analysis reveals long-term accumulation patterns [28-34].

Understanding the long-term accumulation of toxic metals in the human body is essential for several reasons. Firstly, it allows for a more comprehensive evaluation of chronic health effects that might arise from prolonged exposure to these elements [35, 36]. Secondly, this information can inform the development of targeted interventions aimed at reducing exposure to toxic metals and mitigating their associated health risks [37, 38]. Lastly, identifying regional variations in toxic metal accumulation can help address the potential impact of these elements on the health and academic performance of foreign students, who might be more vulnerable to the adverse effects

of toxic metals due to their unique stressors and adaptation challenges [39, 40]. To the best of our knowledge, there are no data on the study of the content of toxic metals: Aluminum (Al), Arsenic (As), Cadmium (Cd), Mercury (Hg), Lead (Pb), and Tin (Sn) in the hair of students from different climatic and geographical regions of the world (Asia, Africa, the Near and Middle East, Latin America) who arrived at the same time in the new identical environmental conditions of the Moscow metropolis. There are only fragmentary studies on the content of individual elements in the literature.

In this study we examined the levels of toxic metals in hair samples of first-year students of RUDN University from various regions, such as Russia, Asia, the Middle East, Africa, and Latin America [41, 42]. The study underscores the importance of comprehending regional differences in toxic metal accumulation and implementing appropriate measures to safeguard the well-being and academic success of foreign students [26, 43].

2. Material and methods

This study was carried out in accordance with the Declaration of Helsinki of the World Medical Association (1964) and its subsequent amendments. All subjects took part in the study voluntarily on the principle of informed consent. The study involved 310 students (40% men, 60% women; mean age 22.7 ± 7.1 years). All subjects were first-year students of the RUDN University. The research involved students from Russia (Moscow and the Moscow region), Southeast Asia (Bangladesh, Vietnam, China, Cambodia, Nepal, Laos, Malaysia, Myanmar, Thailand), Near and Middle East (Iraq, Iran, Yemen, Lebanon, Syria, Palestine, Jordan, Turkey), Africa (Guinea-Bissau, Guinea, Equatorial Guinea, Congo, Ghana, Angola, Zambia, Zimbabwe, Cameroon, Kenya, Ivory Coast, Tanzania, Chad, Benin, Burkina Faso, Rwanda, Sierra Leone) and Latin America (Bolivia, Brazil, Venezuela, Colombia, Mexico, Panama, Peru, Ecuador). The composition of the student cohort by region when entering at RUDN university was as follows: Russia (n = 60), Asia (n = 63), Near and Middle East (n = 81), Africa (n = 55) and Latin America (n = 51). The exclusion criteria were the presence of somatic diseases, as well as specific diets

1 (vegetarianism, raw food diet) and smoking. Immediately after arrival, the subjects were taken hair
2 samples, which is an informative bioindicator substrate in assessing the long-term exposure to
3 toxic metals.
4

5
6 Hair samples were taken from the occipital region using stainless steel scissors pre-treated
7 with ethanol (RUDN University, Moscow, Russia). Only the proximal section of the hair 1–2 cm
8 long was used for analysis (this part of the hair corresponds to the recent time period of the
9 students' life in their region of origin). Subsequently, the proximal parts of the hair strands were
10 washed with acetone, followed by washing three times with distilled deionized water and drying
11 in air. Next, the samples were ground in the presence of 62% nitric acid and placed in a Berghof
12 SpeedWave-4 DAP-40 system (Berghof Products + Instruments GmbH, Germany) for microwave
13 decomposition. Determination of the content of aluminum (Al), arsenic (As), cadmium (Cd),
14 mercury (Hg), lead (Pb) and tin (Sn) in the hair of first-year students was carried out by mass
15 spectrometry with inductively coupled argon (Ar) plasma on a NexION 300D spectrometer
16 (PerkinElmer Inc., USA). The system was calibrated using Universal Data Acquisition Standards
17 Kit (PerkinElmer Inc., USA). Yttrium and rhodium solutions (10 ppb) were used for internal online
18 standardization (Pure Single-Element Standard, PerkinElmer Inc. USA). GBW09101 hair
19 (Shanghai Institute of Nuclear Research, Shanghai, China) was used as a certified reference
20 material of human hair.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 Data processing was performed using the Statistica 10.0 software package (Statsoft, OK,
44 USA). Due to the absence of a Gaussian distribution of data on the content of metals in hair, the
45 median and the corresponding boundaries of the 25–75 centile interval were used as descriptive
46 statistics. The significance of group differences at $p < 0.05$ was assessed using the nonparametric
47 Mann–Whitney U test. For a comparative analysis of the influence of regions (independent
48 predictors) on the level of toxic metals in the hair (dependent variable), we performed a multiple
49 regression analysis adjusted for gender and age of the participants.
50
51
52
53
54
55
56
57
58
59
60

61 **3. Results and discussion**

62
63
64
65

3.1. Analytical figure of merits

Certified reference material of human hair GBW09101 was used for laboratory quality control (**Table 1**). The recovery rates for all studied toxic chemical elements were within the range of 90–110%.

Table 1. Laboratory quality control with certified reference material of human hair GBW09101.

Region	Certified Value, $\mu\text{g/g}$	Obtained Value, $\mu\text{g/g}$	Recovery Rate, %	LoD, ppb	BEC, ppb
Al	23.2	23.3 ± 2.48	101	0.13	0.26
As	0.198	0.194 ± 0.014	97	0.001	0.001
Cd	0.072	0.066 ± 0.006	92	0.0016	0.0006
Hg	1.06	1.15 ± 0.29	108	0.012	0.034
Pb	3.83	4.13 ± 0.28	108	0.0006	0.0007
Sn	n.s.	0.32 ± 0.02	-	n.s.	n.s.

LoD – limit of detection; BEC – background equivalent concentration; ppb – part per billion; n.s. – not specified.

3.2. Hair analysis for toxic metals

The results of the study indicate significant differences in the content of toxic metals in hair samples of first-year students from different regions. For example, the aluminum content (**Table 2**) in hair of African students is 2.4 times higher than that of Russian students, 2.9 times higher than that of Asian students, and 2.5 times higher than that of students from countries of the Nearest and Middle East (NME). Hair samples from students of Latin America also showed higher levels of aluminum compared to students from other countries. Hair samples from Latin American students contain 78% more aluminum than Russian students, 114% more than Asian students, and 88% more than students from the NME region. At the same time, the content of aluminum in the hair of African and Latin American students does not differ significantly.

The content of other elements (arsenic, cadmium, mercury, lead, and tin) in hair samples of first-year students is also characterized by pronounced regional differences (**Tables 2**). For example, the lowest levels of arsenic are observed among Russian students. At the same time, the lowest content of cadmium is typical for representatives of Asian and NME countries, being 2 times lower than the control values. The highest levels of cadmium are found in hair samples from students from Africa and Latin America. The content of cadmium in African students is 4.6 times higher than in Asian students and 5.6 times higher than in NME students. A similar situation is observed in the case of Latin American students (cadmium levels are 4.1 times higher than those for Asian students, and 5 times higher than for students from NME countries). Despite two-fold differences, the level of cadmium in the hair of Russian students is quite high and does not significantly differ from that of African and Latin American students. The content of mercury in hair of students from Latin America exceeds the level of mercury of Russian, Asian and NME students by 118, 75 and 136%, respectively (**Table 2**). The maximum lead content was registered in Africans: their level is higher in comparison with Russian (by 2.7 times), Asian (by 4.5 times) and NME (by 3.8 times) first-year students. Unlike other metals, the content of tin in students' hair is not characterized by significant differences. However, it should be noted that there is a slight trend towards an increase in the level of tin in hair of Latin American and African students.

Table 2. Levels of aluminum, arsenic, cadmium, mercury, lead, and tin in hair ($\mu\text{g/g}$) of foreign students and students from Russia.

Region	Al	As	Cd	Hg	Pb	Sn
Russia	5.59 (3.614÷ 8.443)	0.015 (0.009÷ 0.020)	0.014 (0.006÷ 0.022)	0.104 (0.064÷ 0.164)	0.290 (0.128÷ 0.532)	0.177 (0.06÷ 0.407)
Asia	4.632 (3.544÷	0.037 (0.023÷	0.006 (0.004÷	0.130 (0.064÷	0.173 (0.122÷	0.069 (0.049÷0.2)

	6.663)	0.052) ¹	0.012) ¹	0.172)	0.465)	
NME	5.297	0.030	0.005	0.078	0.207	0.046
	(3.733÷	(0.019÷	(0.003÷	(0.043÷	(0.122÷	(0.025÷
	7.265)	0.049) ¹	0.015) ¹	0.158)	0.38)	0.093)
Africa	13.542	0.039	0.028	0.096	0.794	0.433
	(8.145÷	(0.033÷	(0.006÷	(0.068÷	(0.28÷	(0.16÷
	18.215) ^{1,2,3}	0.057) ¹	0.055) ^{2,3}	0.143) ³	1.826) ^{1,2,3}	0.704)
Latin	9.947	0.038	0.025	0.227	0.435	0.443
America	(5.755÷	(0.026÷	(0.004÷	(0.135÷	(0.113÷	(0.088÷
	12.059) ^{1,2,3}	0.055) ¹	0.097) ^{2,3}	0.456) ^{1,2,3}	1.582)	0.898)

The data are presented as median and 25–75 centile interval; ^{1,2,3} – reliability of differences at $p < 0.05$ relative to the regions: 1 – Russia, 2 – Asia, 3 – NME countries.

We performed multiple linear regression analysis to further analyze the relationship between the region of origin of first-year students and the level of metals in their hair (**Table 3**). We found that geographic differences have the strongest effect on mercury and lead levels in hair, and this is broadly consistent with group comparison results. In particular, living in Asia, Africa, and Latin America is a predictor of high mercury levels in hair. Living in Africa is also significantly associated with higher levels of lead. At the same time, male sex is a negative predictor of the level of metal content in the hair. In the case of cadmium, the direct relationship between residence in Latin America and the level of the metal in hair is close to significant. At the same time, age has a significant effect on the content of cadmium in hair. Between living in different regions and the content of aluminum, arsenic, and tin in hair, we did not reveal any significant relationships using the regression models.

Table 3. Regression analysis of the relationship between the level of metals in hair samples of first-year students and the region of their residence.

Predictor	Al	As	Cd	Hg	Pb	Sn
Male sex	-0.072	0.118	-0.056	-0.224 ($p = 0.001$)	-0.138 ($p = 0.042$)	-0.039
Age, years	-0.001	0.038	0.322 ($p < 0.001$)	-0.030	-0.027	-0.006
Russia	0.005	0.097	0.078	0.153	0.023	0.190
Asia	0.016	0.220	0.074	0.258 ($p = 0.043$)	0.085	0.053
NME	0.118	0.207	-0.007	0.234	0.103	0.153
Africa	0.151	0.182	0.099	0.324 ($p = 0.004$)	0.295 ($p = 0.011$)	0.024
Latin America	0.078	0.106	0.190 ($p = 0.058$)	0.373 ($p < 0.001$)	0.112	0.054

The data are presented as regression coefficients (β), for significant relationships, the corresponding p values are indicated.

Thus, the survey of first-year students at RUDN University shows that foreign students are characterized by higher rates of accumulation of toxic metals in their hair. In particular, the maximum levels of mercury content are observed in students who arrived from Latin America, and the maximum level of lead is typical for African students. Differences in the content of arsenic and cadmium in hair samples of students from different regions are less pronounced. The revealed differences in the patterns of metal accumulation in the body of subjects from different regions most likely reflect characteristic features of environmental pollution by metals in the mentioned countries [1]. High levels of lead in African students are consistent with the ubiquity of lead sources in Africa, while mercury and arsenic emissions are more localized [44]. Hrubá et al. have recently shown that Moroccan blood lead levels are twice as high as those in European countries, China, and Ecuador [38]. This is also consistent with the data obtained in the Ngueta group that children living near the Sahara are characterized by a significant excess of blood lead levels compared to the United States [45]. Lead sources in Africa are primarily gold [46] and lead-zinc [44] mines. The direct routes of lead entry into the human body can be associated with water, soil, grain, and dust pollution [47]. Environmental pollution with cadmium is also an important problem in a number of African countries, including Nigeria [48]. As for arsenic, emissions of this toxicant are not a widespread problem in Africa, and only in some cases, gold mining operations are

1 associated with arsenic emissions into the environment and its subsequent negative impact on
2 human health [49].
3

4 Despite the leading role of Asian countries (primarily China) in the release of mercury into
5 the atmosphere, the maximum content of mercury in hair is characteristic of students from Latin
6 America. This observation is consistent with previously found by Hrubá et al. high concentrations
7 of mercury and lead in Ecuador [38]. The leading role of mercury as an inorganic pollutant in the
8 Latin American region may be due to the prevalence of artisanal gold mining [50]. For example,
9 Streets et al. state that the direct contribution of Latin America to the total release of mercury into
10 the atmosphere is 10% [51]. Along with mercury, environmental pollution with arsenic is also an
11 urgent medical and social problem in the Latin American region, and it is associated with high
12 incidence of pathology caused by arsenic exposure [52].
13
14
15
16
17
18
19
20
21
22
23
24
25

26 Living in Asia is also a significant predictor of metal accumulation in the body, associated
27 with increased concentrations of mercury and cadmium. This observation is consistent with the
28 recent work of Dix-Cooper and Kosatsky, who reported higher concentrations of mercury, lead
29 and cadmium in whole blood in women that arrived from Asia in Canada compared with the
30 Canadian population [26]. Some of us also recently conducted a survey of residents of the Republic
31 of Bangladesh and found higher levels of mercury and arsenic in hair compared to Russian ones
32 [53]. Moreover, the reported concentrations of toxic metals in Asian first-year students are in close
33 agreement with those of Guangzhou college students published by Zhang et al. [54]. Despite
34 significant preventive activities, metal pollution remains a significant problem in Central and
35 Southeast Asia [55]. The risk of excessive exposure to mercury also remains for residents of the
36 Near and Middle East, and this is consistent with the observed annual increase in mercury
37 emissions in the region during the period from 2000 to 2015 [51]. The development of the
38 metalworking industry in the Middle East also makes a significant contribution to environmental
39 pollution with cadmium and lead [56]. Considering the identified stress reactions in first-year
40 foreign students of RUDN University [57], excessive accumulation of metals due to environmental
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 pollution of the regions from which these students came might significantly reduce the adaptive
2 reactions of the students through the implementation of a wide range of toxicity mechanisms [58,
3 59]. This negative effect, in turn, might contribute to an increase in illness and a decrease in the
4 effectiveness of teaching foreign students and should be the subject of future studies.
5
6

7
8
9 Our findings revealed increased accumulation of different toxic metals in students hailing
10 from different geographic regions. It seems natural that the detected increased accumulation
11 mirrors the specific features of environmental pollution with metals in these countries, which can
12 result from factors such as industrial activities, mining, and inadequate waste management
13 practices [60, 61].
14
15
16
17
18
19
20

21 **4. Conclusion**

22 The results of this work can be considered from the following few perspectives.
23
24

25
26
27
28 Firstly, we found an increased accumulation of toxic metals, especially mercury, lead,
29 cadmium, in the body of first-year students, who arrived primarily from Africa and Latin America.
30
31
32

33
34 Secondly, we confirmed that the differences in the patterns of metal accumulation in the
35 students from different regions most likely reflect the specific features of environmental pollution
36 with metals in these countries. Probably, this increased accumulation of toxic metals might have a
37 negative impact on the health status and effectiveness of teaching foreign students. Therefore,
38 further research is needed, focused on assessing the direct contribution of excess toxic metals to
39 the health indicators of RUDN University students, and this project is underway in our group.
40
41
42
43
44
45
46
47

48 **Acknowledgements**

49
50
51
52 This paper has been supported by the RUDN University Strategic Academic Leadership
53 Program (recipient: A.A. Kirichuk, award no. 202713-0-000 “Development of a scientifically
54 based methodology for the ecological adaptation of foreign students to new environmental
55 conditions”).
56
57
58
59
60
61
62
63
64
65

References

- 1 [1] Y. Li, Q. Zhou, B. Ren, J. Luo, J. Yuan, X. Ding, H. Bian, X. Yao, Trends and Health Risks
2 of Dissolved Heavy Metal Pollution in Global River and Lake Water from 1970 to 2017, in: P. de
3 Voogt (Ed.), Reviews of Environmental Contamination and Toxicology Volume 251, Springer
4 International Publishing, Cham, 2020, pp. 1-24.
5
6
7
8
9
10
11 [2] F. Cusset, S.J. Reynolds, A. Carravieri, D. Amouroux, O. Asensio, R.C. Dickey, J. Fort, B.J.
12 Hughes, V.H. Paiva, J.A. Ramos, L. Shearer, E. Tessier, C.P. Wearn, Y. Cherel, P. Bustamante, A
13 century of mercury: Ecosystem-wide changes drive increasing contamination of a tropical seabird
14 species in the South Atlantic Ocean, *Environ. Pollut.* 323 (2023) 121187.
15
16
17
18
19
20
21 <https://doi.org/10.1016/j.envpol.2023.121187>.
22
23
24 [3] L.P. Padhye, T. Jasemizad, S. Bolan, O.V. Tsyusko, J.M. Unrine, B.K. Biswal, R.
25 Balasubramanian, Y. Zhang, T. Zhang, J. Zhao, Y. Li, J. Rinklebe, H. Wang, K.H.M. Siddique,
26 N. Bolan, Silver contamination and its toxicity and risk management in terrestrial and aquatic
27 ecosystems, *Sci. Total Environ.* 871 (2023) 161926.
28
29
30
31
32
33 <https://doi.org/10.1016/j.scitotenv.2023.161926>.
34
35
36 [4] Y. Zhou, Y. Tang, C. Liao, M. Su, K. Shih, Recent advances toward structural incorporation
37 for stabilizing heavy metal contaminants: A critical review, *J. Hazard. Mater.* 448 (2023) 130977.
38
39
40
41
42 <https://doi.org/10.1016/j.jhazmat.2023.130977>.
43
44 [5] A. Kakade, M. Sharma, E.-S. Salama, P. Zhang, L. Zhang, X. Xing, J. Yue, Z. Song, L. Nan,
45 S. Yujun, X. Li, Heavy metals (HMs) pollution in the aquatic environment: Role of probiotics and
46 gut microbiota in HMs remediation, *Environ. Res.* 223 (2023) 115186.
47
48
49
50
51
52 <https://doi.org/10.1016/j.envres.2022.115186>.
53
54 [6] H. Khawashi, V. Chavan, D. Bhagat, S. Deshmukh, Recent Advances in the Detection of Lead
55 Ions using Nanoparticle-Based Sensors, *Biointerface Res. Appl. Chem.* 13 (2023) 466.
56
57
58
59
60
61
62
63
64
65 <https://doi.org/10.33263/BRIAC135.466>.

1 [7] A. Pandey, A. Kalamdhad, Y. Chandra Sharma, Recent advances of nanocellulose as biobased
2 adsorbent for heavy metal ions removal: A sustainable approach integrating with waste
3 management, *Environ. Nanotechnol. Monit. Manag.* 20 (2023) 100791.

4 <https://doi.org/10.1016/j.enmm.2023.100791>.

5
6
7
8 [8] P.B. Tchounwou, C.G. Yedjou, A.K. Patlolla, D.J. Sutton, Heavy Metal Toxicity and the
9 Environment, in: A. Luch (Ed.), *Molecular, Clinical and Environmental Toxicology: Volume 3:*
10 *Environmental Toxicology*, Springer Basel, Basel, 2012, pp. 133-164.

11
12
13 [9] A. Manouchehri, S. Shokri, M. Pirhadi, M. Karimi, S. Abbaszadeh, G. Mirzaei, M. Bahmani,
14 The Effects of Toxic Heavy Metals Lead, Cadmium and Copper on the Epidemiology of Male and
15 Female Infertility, *JBRA Assist. Reprod.* 26(4) (2022) 627-630. <https://doi.org/10.5935/1518->
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66 [10] S. Dutta, B. Gorain, H. Choudhury, S. Roychoudhury, P. Sengupta, Environmental and
67 occupational exposure of metals and female reproductive health, *Environ. Sci. Pollut. Res.* 29(41)
68 (2022) 62067-62092. <https://doi.org/10.1007/s11356-021-16581-9>.

69 [11] R.L. Imhof, D.M.R. Davis, M.M. Tollefson, Hair Loss, *Pediatr. Rev.* 41(11) (2020) 570-584.
70 <https://doi.org/10.1542/pir.2019-0009>.

71 [12] C. Yang, Z. Wang, The Epitranscriptomic Mechanism of Metal Toxicity and Carcinogenesis,
72 *Int. J. Mol. Sci.* 23(19) (2022) 11830.

73 [13] H. Goel, K. Goyal, A.K. Pandey, M. Benjamin, F. Khan, P. Pandey, S. Mittan, D. Iqbal, M.
74 Alsaweed, W. Alturaiki, Y. Madkhali, M.A. Kamal, P. Tanwar, T.K. Upadhyay, Elucidations of
75 Molecular Mechanism and Mechanistic Effects of Environmental Toxicants in Neurological
76 Disorders, *CNS Neurol. Disord. Drug Targets* 22(1) (2023) 84-97.
77 <https://doi.org/10.2174/1871527321666220329103610>.

78 [14] A. Shiani, K. Sharafi, A.K. Omer, A. Kiani, B. Karamimatin, T. Massahi, G. Ebrahimzadeh,
79 A systematic literature review on the association between exposures to toxic elements and an

- autism spectrum disorder, *Sci. Total Environ.* 857 (2023) 159246.
<https://doi.org/10.1016/j.scitotenv.2022.159246>.
- [15] S.M. Awadh, Z.M. Yaseen, M.S. Al-Suwaiyan, The role of environmental trace element toxicants on autism: A medical biogeochemistry perspective, *Ecotoxicol. Environ. Saf.* 251 (2023) 114561. <https://doi.org/10.1016/j.ecoenv.2023.114561>.
- [16] M. Taghavi, M. Darvishiyan, M. Momeni, H. Eslami, R.A. Fallahzadeh, A. Zarei, Ecological risk assessment of trace elements (TEs) pollution and human health risk exposure in agricultural soils used for saffron cultivation, *Sci. Rep.* 13(1) (2023) 4556. <https://doi.org/10.1038/s41598-023-31681-x>.
- [17] H. Ali, E. Khan, I. Ilahi, Environmental Chemistry and Ecotoxicology of Hazardous Heavy Metals: Environmental Persistence, Toxicity, and Bioaccumulation, *J. Chem.* 2019 (2019) 6730305. <https://doi.org/10.1155/2019/6730305>.
- [18] Z. Li, Z. Ma, T.J. van der Kuijp, Z. Yuan, L. Huang, A review of soil heavy metal pollution from mines in China: Pollution and health risk assessment, *Sci. Total Environ.* 468-469 (2014) 843-853. <https://doi.org/10.1016/j.scitotenv.2013.08.090>.
- [19] G. Ngueta, R. Ndjaboue, Blood lead concentrations in sub-Saharan African children below 6 years: systematic review, *Trop. Med. Int. Health* 18(10) (2013) 1283-91. <https://doi.org/10.1111/tmi.12179>.
- [20] S. Karatela, C. Coomarasamy, J. Paterson, N.I. Ward, Exposure to Toenail Heavy Metals and Child Behavior Problems in Nine-Year-Old Children: A Cross-Sectional Study, *Int. J. Environ. Res. Public Health* 17(11) (2020) 4120.
- [21] A. Błażewicz, E. Grywalska, P. Macek, P. Mertowska, S. Mertowski, J. Wojnicka, N. Durante, A. Makarewicz, Research into the Association of Cadmium and Manganese Excretion with Thyroid Function and Behavioral Areas in Adolescents with Autism Spectrum Disorders, *J. Clin. Med.* 11(3) (2022) 579.

- 1 [22] L. Malavika, T. Goyal, P. Mitra, G. Saikiran, S. Sharma, P. Sharma, Risk Factors for Lead
2 Toxicity and its Effect on Neurobehavior in Indian Children, *Indian J. Clin. Biochem.* 37(3) (2022)
3 294-302. <https://doi.org/10.1007/s12291-021-00995-w>.
4
5 [23] Y. Qin, C. Xu, W. Li, B. Jian, B. Wu, M. Chen, H. Sun, H. Hong, Metal/metalloid levels in
6 hair of Shenzhen residents and the associated influencing factors, *Ecotoxicol. Environ. Saf.* 220
7 (2021) 112375. <https://doi.org/10.1016/j.ecoenv.2021.112375>.
8
9 [24] R. Canham, A.M. González-Prieto, J.E. Elliott, Mercury Exposure and Toxicological
10 Consequences in Fish and Fish-Eating Wildlife from Anthropogenic Activity in Latin America,
11 *Integr. Environ. Assess. Manag.* 17(1) (2021) 13-26. <https://doi.org/10.1002/ieam.4313>.
12
13 [25] L. Lv, B. Liu, Y. Yu, W. Dong, L. Gao, Y. He, Heavy metals in paired samples of hair and
14 nails in China: occurrence, sources and health risk assessment, *Environ. Geochem. Health* (2022).
15
16 <https://doi.org/10.1007/s10653-022-01400-y>.
17
18 [26] L. Dix-Cooper, T. Kosatsky, Blood mercury, lead and cadmium levels and determinants of
19 exposure among newcomer South and East Asian women of reproductive age living in Vancouver,
20 Canada, *Sci. Total Environ.* 619-620 (2018) 1409-1419.
21
22 <https://doi.org/10.1016/j.scitotenv.2017.11.126>.
23
24 [27] J. Zheng, M. Li, B. Tang, W. Luo, Y. Ma, M. Ren, Y. Yu, X. Luo, B. Mai, Levels, Spatial
25 Distribution, and Impact Factors of Heavy Metals in the Hair of Metropolitan Residents in China
26 and Human Health Implications, *Environ. Sci. Technol.* 55(15) (2021) 10578-10588.
27
28 <https://doi.org/10.1021/acs.est.1c02001>.
29
30 [28] T.-H. Ying, C.-J. Huang, C.-J. Hsieh, P.-J. Wu, C.-C. Yeh, P.-K. Hung, W.-H. Chang, M.-H.
31 Wu, H. Hung, J.-W. Chang, C.-T. Wang, R.D. Arcega, T. Mahmudiono, C.-C. Lee, H.-L. Chen,
32 Potential Factors Associated with the Blood Metal Concentrations of Reproductive-Age Women
33 in Taiwan, *Expos. Health* (2023). <https://doi.org/10.1007/s12403-023-00541-4>.
34
35 [29] S. Khan, T.G. Kazi, J.A. Baig, H.I. Afridi, N.F. Kolachi, Separation/Preconcentration
36 Methods for the Determination of Aluminum in Dialysate Solution and Scalp Hair Samples of
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- Kidney Failure Patients, *Biological trace element research* 144(1) (2011) 205-216.
<https://doi.org/10.1007/s12011-011-9070-5>.
- [30] A.Q. Shah, T.G. Kazi, J.A. Baig, H.I. Afridi, Correlation Between Arsenic Concentration in Fish and Human Scalp Hair of People Living in Arsenic-Contaminated and Noncontaminated Areas of Pakistan, *Biological trace element research* 144(1) (2011) 197-204.
<https://doi.org/10.1007/s12011-011-9067-0>.
- [31] J.A. Baig, T.G. Kazi, M.B. Arain, H.I. Afridi, K.P. Mahar, Samples of Kidney Failure Patients by Cloud Point Extraction and Electrothermal Atomic Absorption Spectroscopy, *Journal of AOAC INTERNATIONAL* 95(6) (2019) 1755-1760. <https://doi.org/10.5740/jaoacint.11-303>.
- [32] S. Baloch, T.G. Kazi, H.I. Afridi, J.A. Baig, F.N. Talpur, M.B. Arain, Correlation of lithium levels between drinking water obtained from different sources and scalp hair samples of adult male subjects, *Environmental geochemistry and health* 39(5) (2017) 1191-1199.
<https://doi.org/10.1007/s10653-016-9886-1>.
- [33] S. Baloch, T.G. Kazi, J.A. Baig, H.I. Afridi, M.B. Arain, Occupational exposure of lead and cadmium on adolescent and adult workers of battery recycling and welding workshops: Adverse impact on health, *Science of The Total Environment* 720 (2020) 137549.
<https://doi.org/10.1016/j.scitotenv.2020.137549>.
- [34] K. Rajput, H.I. Afridi, T.G. Kazi, F.N. Talpur, J.A. Baig, Sodium, Potassium, Calcium, and Magnesium in the Scalp Hair and Blood Samples Related to the Clinical Stages of the Parkinson's Disease, *Biological trace element research* 199(7) (2021) 2582-2589.
<https://doi.org/10.1007/s12011-020-02399-y>.
- [35] J. Zheng, M. Li, B. Tang, W. Luo, Y. Ma, M. Ren, Y. Yu, X. Luo, B. Mai, Levels, Spatial Distribution, and Impact Factors of Heavy Metals in the Hair of Metropolitan Residents in China and Human Health Implications, *Environ. Sci. Technol.* 55(15) (2021) 10578-10588.
<https://doi.org/10.1021/acs.est.1c02001>.

- 1 [36] J. Webb, N. Mainville, D. Mergler, M. Lucotte, O. Betancourt, R. Davidson, E. Cueva, E.
2 Quizhpe, Mercury in Fish-eating Communities of the Andean Amazon, Napo River Valley,
3 Ecuador, *EcoHealth* 1 (2004) SU59-SU71. <https://doi.org/10.1007/s10393-004-0063-0>.
4
5 [37] E. Blaurock-Busch, O.R. Amin, H.H. Dessoki, T. Rabah, Toxic Metals and Essential
6 Elements in Hair and Severity of Symptoms among Children with Autism, *Maedica* 7(1) (2012)
7 38-48.
8
9 [38] F. Hrubá, U. Strömberg, M. Černá, C. Chen, F. Harari, R. Harari, M. Horvat, K. Koppová, A.
10 Kos, A. Krsková, M. Krsnik, J. Laamech, Y.F. Li, L. Löfmark, T. Lundh, N.G. Lundström, B.
11 Lyoussi, D. Mazej, J. Osredkar, K. Pawlas, N. Pawlas, A. Prokopowicz, G. Rentschler, V.
12 Spěváčková, Z. Spiric, J. Tratnik, S. Skerfving, I.A. Bergdahl, Blood cadmium, mercury, and lead
13 in children: an international comparison of cities in six European countries, and China, Ecuador,
14 and Morocco, *Environ. Int.* 41 (2012) 29-34. <https://doi.org/10.1016/j.envint.2011.12.001>.
15
16 [39] E. Brodtkin, R. Copes, A. Mattman, J. Kennedy, R. Kling, A. Yassi, Lead and mercury
17 exposures: interpretation and action, *Can. Med. Assoc. J.* 176(1) (2007) 59-63.
18 <https://doi.org/10.1503/cmaj.060790>.
19
20 [40] A. Vienna, E. Capucci, M. Wolfsperger, G. Hauser, Heavy metal concentration in hair of
21 students in Rome, *Anthropol. Anz.* 53(1) (1995) 27-32.
22
23 [41] A. Monger, K. Wangdi, Lead and Mercury Exposure and Related Health Problems in Metal
24 Artisan Workplaces and High-Risk Household Contacts in Thimphu, Bhutan, *Sci. World J.* 2020
25 (2020) 9267181. <https://doi.org/10.1155/2020/9267181>.
26
27 [42] A. Wilk, M. Romanowski, B. Wiszniewska, Analysis of Cadmium, Mercury, and Lead
28 Concentrations in Erythrocytes of Renal Transplant Recipients from Northwestern Poland,
29 *Biology* 10(1) (2021). <https://doi.org/10.3390/biology10010062>.
30
31 [43] M. Szyrkowska, A. Pawlaczyk, M. Elżbieta, S. Sypniewski, T. Paryjczak, Human Hair as a
32 biomarker in assessing exposure to toxic metals, *Pol. J. Environ. Stud.* 18 (2009) 1151-1161.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [44] J. Yabe, M. Ishizuka, T. Umemura, Current levels of heavy metal pollution in Africa, *J. Vet. Med. Sci.* 72(10) (2010) 1257-1263. <https://doi.org/10.1292/jvms.10-0058>.
- [45] G. Ngueta, R. Ndjaboue, Blood lead concentrations in sub-Saharan African children below 6 years: systematic review, *Trop. Med. Int. Health* 18(10) (2013) 1283-1291. <https://doi.org/10.1111/tmi.12179>.
- [46] C.A. Dooyema, A. Neri, Y.-C. Lo, J. Durant, P.I. Dargan, T. Swarthout, O. Biya, S.O. Gidado, S. Haladu, N. Sani-Gwarzo, P.M. Nguku, H. Akpan, S.a. Idris, A.M. Bashir, M.J. Brown, Outbreak of Fatal Childhood Lead Poisoning Related to Artisanal Gold Mining in Northwestern Nigeria, 2010, *Environ. Health Perspect.* 120(4) (2012) 601-607. <https://doi.org/10.1289/ehp.1103965>.
- [47] O. Bello, R. Naidu, M.M. Rahman, Y. Liu, Z. Dong, Lead concentration in the blood of the general population living near a lead–zinc mine site, Nigeria: Exposure pathways, *Sci. Total Environ.* 542 (2016) 908-914. <https://doi.org/10.1016/j.scitotenv.2015.10.143>.
- [48] O.E. Orisakwe, Lead and cadmium in public health in Nigeria: physicians neglect and pitfall in patient management, *N. Am. J. Med. Sci.* 6(2) (2014) 61-70. <https://doi.org/10.4103/1947-2714.127740>.
- [49] C. Kamunda, M. Mathuthu, M. Madhuku, Health Risk Assessment of Heavy Metals in Soils from Witwatersrand Gold Mining Basin, South Africa, *Int. J. Environ. Res. Public Health* 13(7) (2016) 663.
- [50] J. Drewry, J. Shandro, M.S. Winkler, The extractive industry in Latin America and the Caribbean: health impact assessment as an opportunity for the health authority, *Int. J. Public Health* 62(2) (2017) 253-262. <https://doi.org/10.1007/s00038-016-0860-6>.
- [51] D.G. Streets, H.M. Horowitz, Z. Lu, L. Levin, C.P. Thackray, E.M. Sunderland, Global and regional trends in mercury emissions and concentrations, 2010–2015, *Atmos. Environ.* 201 (2019) 417-427. <https://doi.org/10.1016/j.atmosenv.2018.12.031>.

- 1 [52] K.M. Khan, R. Chakraborty, J. Bundschuh, P. Bhattacharya, F. Parvez, Health effects of
2 arsenic exposure in Latin America: An overview of the past eight years of research, *Sci. Total*
3 *Environ.* 710 (2020) 136071. <https://doi.org/10.1016/j.scitotenv.2019.136071>.
4
5 [53] O.A. Skalnaya, A.A. Skalnaya, V.A. Demidov, N.G. Zhivaev, Hair mineral element content
6 in adult inhabitants of Mimensingh district, Bangladesh republic. Report 1: Toxic elements (As,
7 Be, Cd, Hg, Pb), *Mikroelementy v meditsine* 16(3) (2015) 45-49.
8
9 [54] S. Lu, L. Ren, J. Fang, J. Ji, G. Liu, J. Zhang, H. Zhang, R. Luo, K. Lin, R. Fan, Trace elements
10 are associated with urinary 8-hydroxy-2'-deoxyguanosine level: a case study of college students
11 in Guangzhou, China, *Environ. Sci. Pollut. Res.* 23(9) (2016) 8484-8491.
12 <https://doi.org/10.1007/s11356-016-6104-8>.
13
14 [55] Z. Barnett-Itzhaki, M. Esteban López, N. Puttaswamy, T. Berman, A review of human
15 biomonitoring in selected Southeast Asian countries, *Environ. Int.* 116 (2018) 156-164.
16 <https://doi.org/10.1016/j.envint.2018.03.046>.
17
18 [56] N. Sistani, M. Moeinaddini, N. Khorasani, A. Hamidian, M. Ali-Taleshi, R. Azimi
19 Yancheshmeh, Heavy metal pollution in soils nearby Kerman steel industry: metal richness and
20 degree of contamination assessment, *Iran. J. Health Saf. Environ.* 10(1) (2017) 75-86.
21
22 [57] A.A. Kirichuk, Stress reaction and reactivity of the cardiovascular system of foreign students
23 of the Peoples' Friendship University of Russia, *Technol. Living Syst.* 17(4) (2020) 71-77.
24
25 [58] M. Jaishankar, T. Tseten, N. Anbalagan, B.B. Mathew, K.N. Beeregowda, Toxicity,
26 mechanism and health effects of some heavy metals, *Interdiscip. Toxicol.* 7(2) (2014) 60-72.
27 <https://doi.org/10.2478/intox-2014-0009>.
28
29 [59] A.A. Kirichuk, A.A. Skalny, A.I. Rusakov, A.A. Tinkov, A.V. Skalny, Arsenic, cadmium,
30 mercury, and lead levels in hair and urine in first-year RUDN University students of different
31 geographic origins, *Environ. Sci. Pollut. Res.* 27(27) (2020) 34348-34356.
32 <https://doi.org/10.1007/s11356-020-09683-3>.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 [60] M.C.O. Souza, B.A. Rocha, J.A. Adeyemi, M. Nadal, J.L. Domingo, F. Barbosa, Legacy and
2 emerging pollutants in Latin America: A critical review of occurrence and levels in environmental
3 and food samples, *Sci. Total Environ.* 848 (2022) 157774.
4
5 <https://doi.org/10.1016/j.scitotenv.2022.157774>.
6

7
8 [61] Y. Irawati, H. Kusnopranto, U.F. Achmadi, A. Safrudin, A. Sitorus, R. Risandi, S.
9 Wangsamuda, P.B. Setia Asih, D. Syafruddin, Blood lead levels and lead toxicity in children aged
10 1-5 years of Cinangka Village, Bogor Regency, *PLoS One* 17(2) (2022) e0264209.
11
12 <https://doi.org/10.1371/journal.pone.0264209>.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65