



Thermophysical properties of H₂/gas/brine system under subsurface storage conditions: molecular simulations and thermodynamic modeling

Halla Kerkache, Hai Hoang, Pierre Cézac, Salaheddine Chabab, Guillaume Galliero

► To cite this version:

Halla Kerkache, Hai Hoang, Pierre Cézac, Salaheddine Chabab, Guillaume Galliero. Thermophysical properties of H₂/gas/brine system under subsurface storage conditions: molecular simulations and thermodynamic modeling. American Institute of Chemical Engineers (AIChE), Nov 2023, Orlando (FL), United States. hal-04624333

HAL Id: hal-04624333

<https://univ-pau.hal.science/hal-04624333>

Submitted on 25 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Thermophysical properties of H₂/gas/brine system under subsurface storage conditions: molecular simulations and thermodynamic modeling



Halla Kerkache^{a,b}, Hai Hoang^a, Pierre Cézac^b, Salaheddine Chabab^{a,b}, Guillaume Galliéro^a


^a Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, TOTALENERGIES, LFCR, UMR5150, Pau, France
^b Université de Pau et des Pays de l'Adour, E2S UPPA, LaTEP, Pau, France



Context & Motivations

The storage of hydrogen in deep saline aquifers offer a great capacity and geographical availability for large-scale renewable energy storage. However, the contact between gas, brine, and rock can affect the mobility and stability of hydrogen and is not sufficiently documented at the time being. In the first part of this work, by using Monte Carlo simulation, new equilibrium data for H₂/brine systems are generated. Then, by using non-equilibrium molecular dynamics, new data on self-diffusivity for H₂/NaCl brine systems are predicted for a wide range of conditions relevant to underground hydrogen storage in porous media. Finally, the co-solubility of gases in the ternary systems CO₂-CH₄-H₂O and CO₂-H₂-H₂O was modeled. An example of the application of this model is given by the prediction of optimal conditions (CO₂/H₂ ratio) at equilibrium to ensure the stoichiometry of the in-situ methanation reaction.

Selected Force Fields



Monte Carlo Simulation details

Sampling method: Continuous Fractional Component Monte Carlo (CFCMC)/NPT ensemble

Henry constant from molecular simulation

$$H_i = \rho k_B T \exp(\mu_i^{\text{excess}} / k_B T)$$

Molar composition

$$x_i(P, T, m_s) = \frac{y_i P \varphi_i}{H_i \gamma_i \exp\left(\vartheta_i^\infty \frac{P - P_{\text{sat}, \text{solvent}}}{RT}\right)}$$

Dynamics Simulation details


Method: External field non-equilibrium molecular dynamics (EF-NEMD).

Estimation of the Self-diffusion coefficient (SDC)


$$D_{\text{Gas}}^{\text{NEMD}} = \frac{\mu_{\text{Gas}} RT}{F_{\text{Gas}}^{\text{Ext}}}$$

Monte Carlo Simulation


Henry constant of H₂ in NaCl brine



Radial distribution function



Molar composition of H₂ in NaCl brine



- ✓ Excellent representation of H₂ solubility in NaCl brine using three different models (Marx_TIP4P/2005_OPLS – Marx_TIP4P/EP_KBF-Marx_TIP4P/2005_Madrid).
- ✓ The radial distribution function (RDF) is used to evaluate the solvation structure of the H₂/brine mixtures which is shown to be dominated by the H₂-H₂O and H₂-Cl⁻ interactions. A constant BIP on energy for H₂-H₂O and on volume for H₂-Cl⁻ are used to improve the quantification of the H₂ solubility in brine.

Non-Equilibrium molecular dynamics

H ₂ -H ₂ O	Approaches	T [K]	P [bar]	D × 10 ⁹ [m ² /s]	σ
Marx + TIP4P/2005 [2]	EMD	298.15	1	5.90	1.8
Vrabec + TIP4P/2005 [3]	EMD	298.15	1	4.40	0.6
Marx + TIP4P/2005 –This work	NEMD	298.15	1	5.20	0.28
H ₂ -H ₂ O-NaCl (m _{NaCl} =1 mol _{NaCl} /kg _w)	Approaches	T [K]	P [bar]	D × 10 ⁹ [m ² /s]	σ
Vrabec + TIP4P/2005+Madrid Transport [3]	EMD	298.15	1	3.70	0.6
Marx + TIP4P/2005 +Madrid –This work	NEMD	298.15	1	4.14	0.13
Marx + TIP4P/2005 +Madrid Transport –This work	NEMD	298.15	1	4.45	0.22

- ✓ The predicted self-diffusion coefficients for H₂ in pure water and NaCl brine are within the order of magnitude and very similar to literature results, with a better (lower) uncertainty.


- ✓ H₂ Diffusion tends to decrease in brine relatively to pure water

EMD: Equilibrium molecular dynamics

NEMD: Non-Equilibrium molecular dynamics



Thermodynamic Modeling



- ✓ The parameters of both models are fitted on binary (gas/water) solubility data.
- ✓ The models predict the ternary CO₂-CH₄-H₂O system very well when compared with literature experimental data.
- ✓ The ternary H₂-CO₂-H₂O system was also modeled to estimate the optimal H₂/CO₂ ratio for underground bio-methanation application.



Conclusions & Perspectives

- Molecular simulation models were validated and used to provide "pseudo-experimental" data. These data in addition to those from the literature for other gases (CH₄ and CO₂) were used to optimize the parameters of the thermodynamic models (phi-phi and gamma-phi approaches).
- The thermodynamic modeling showed that the optimal H₂/CO₂ ratio for underground bio-methanation application is around 3% CO₂ (97% H₂) to be injected (to have a stoichiometric ratio of 1:4).
- A good representation of H₂ self-diffusion coefficient in water and brine are obtained using External Forces Non-equilibrium Molecular Dynamics.
- The possible influence of the clay interlayer on solubilities and diffusivities will be explored.

References

