



HAL
open science

Comparative study of three H₂ geological storages in deep aquifers simulated in high pressure reactors.

Jean Mura, Magali Ranchou-Peyruse, Marion Guignard, Perla Haddad, Marion Ducouso, Franck Casteran, Pascale Sénéchal, Marie-Pierre Isaure, Peter Moonen, Guilhem Hoareau, et al.

► **To cite this version:**

Jean Mura, Magali Ranchou-Peyruse, Marion Guignard, Perla Haddad, Marion Ducouso, et al.. Comparative study of three H₂ geological storages in deep aquifers simulated in high pressure reactors.. Goldschmidt 2023, Jul 2023, Lyon, France. <10.7185/gold2023.18303>. <hal-05325790>

HAL Id: hal-05325790

<https://univ-pau.hal.science/hal-05325790v1>

Submitted on 22 Oct 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Comparative study of three H₂ geological storages in deep aquifers simulated in high pressure reactors.

JEAN MURA¹, MAGALI RANCHOPEYRUSE^{1,2,3},
MARION GUIGNARD³, PERLA HADDAD¹, MARION
DUCOUSSO^{1,2}, FRANCK CASTERAN^{1,2}, PASCALE
SÉNÉCHAL⁴, MARIE-PIERRE ISAURE³, PETER
MOONEN⁴, GUILHEM HOAREAU⁵, ALICE BALDY³,
ANÉLIA PETIT⁶, PIERRE CHIQUET^{2,7}, GUILHEM
CAUMETTE^{2,8}, PIERRE CÉZAC^{1,2} AND ANTHONY
RANCHOPEYRUSE^{2,3}

¹Université de Pau et des Pays de l'Adour, E2S UPPA, LaTEP

²Joint Laboratory SEnGA, UPPA-E2S-Teréga

³Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM

⁴Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, DMEX

⁵Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, TOTAL, LFCR

⁶STORENGY – Geosciences Department

⁷Teréga – Geosciences Department

⁸Teréga – Environment Department

Presenting Author: jean.mura@univ-pau.fr

In the context of climate change and resource depletion, an adaptation in the energy mix towards decarbonation and renewable energy is crucial. Dihydrogen (H₂) is a promising alternative to traditional carbonated energy sources. Besides being storable, it also has the potential to be produced using renewable and low carbon processes. In order to use H₂ on a large scale, it will be necessary to store massive quantities by means of, for example, Underground Gas Storage (UGS) in deep aquifers.

H₂'s behavior in deep aquifer is related to its geochemical reactivity and to the microbial activity. Also, it is an electron donor as well as an energy source for numerous indigenous microorganisms. In this study, H₂ injection in three different UGS, with different formation waters, rocks and microbial communities, were simulated in a high-pressure reactor following a previously defined protocol^[1].

To better understand the intricate phenomena at work, extent of reaction equations based on microbial diversities were solved to identify the main reactions taking place in the reactor. The broadly used geochemical modeling software PHREEQC was used to calculate gases solubilities, resulting pH and redox potential inside the reactor.

While methanogens were only observed in some of the experiments, the appearance of formate seems to be a constant and should therefore be considered in the evolution of storage conditions. The effect of H₂ on microbial communities and thus their activities seem to extend well beyond the metabolisms usually targeted for hydrogenotrophy.

In the first experiment with low sulfate concentration, sulfate

was totally depleted during the experiment and H₂ consumption stopped when CO₂ had disappeared. In the two other experiments with higher sulfate concentration, H₂ consumption was observed throughout the experiment while CO₂ and sulfate remained at the end of experiment. Acetogenesis was also observed on a lower scale. It appeared that H₂ injection led to alkalization, mainly through the consumption of CO₂ by sulfate-reducers and methanogens micro-organisms. Depending on the UGS conditions, especially sulfate concentration, the microbial reactions and their scale varied, highlighting the importance of a site-specific study.

[1] Haddad et al. 2022, Energy Environ. Sci. 15, 3400.